

The graviton vacuum as a distributional state in kinematic Loop Quantum Gravity

Madhavan Varadarajan^{1,2,*}

¹*Raman Research Institute, Bangalore 560 080, India*

²*Sektion Physik, LMU, Theresienstr.37, Munich, Germany*

Abstract

The quantum behaviour of weak gravitational fields admits an adequate, albeit approximate, description by those graviton states in which the expectation values and fluctuations of the linearised gravitational field are small. Such states must approximate corresponding states in full quantum gravity. We analyse the nature of this approximation for the graviton vacuum state in the context of kinematical Loop Quantum Gravity (LQG) wherein the constraints are ignored. We identify the graviton vacuum state with kinematically non-normalizable, distributional states in LQG by demanding that relations between linearised operator actions on the former are mirrored by those of their non-linear counterparts on the latter. We define a semi-norm on the space of kinematical distributions and show that the identification is approximate upto distributions which are small in this semi-norm. We argue that our candidate states are annihilated by the linearised constraints (expressed as operators in the full theory) to leading order in the parameter characterising the approximation. This suggests the possibility, in a scheme such as ours, of solving the full constraints order by order in this parameter. The main drawback of our considerations is that they depend on certain auxiliary constructions which, though mathematically well defined, do not arise from physical insight. Our work is an attempt to implement an earlier proposal of Iwasaki and Rovelli.

PACS numbers:

*Electronic address: madhavan@rri.res.in

I. INTRODUCTION

A Dirac constraint quantization of a Hamiltonian formulation of gravity is defined through the following steps. First, a “kinematical” representation of the Poisson bracket algebra of a large enough set of functions on the unconstrained phase space is constructed such that these functions act as linear operators on the representation space. Next, the constraints of the theory are represented as quantum operators in this representation and physical states are identified with their kernel. Finally, an inner product on the space of physical states is chosen which enforces hermiticity conditions on a complete set of operators corresponding to Dirac observables, thus converting the physical state space to a Hilbert space. Since one expects the Dirac observables to have direct physical interpretation, the physical interpretation of the formalism follows.¹ Within the Loop Quantum Gravity (LQG) approach, the kinematical structures are well understood. There has also been significant progress in finding physical states. Further progress is obstructed by the fact that almost no Dirac observables are known for general relativity. Indeed, this is a problem of the *classical* theory. As in any generally covariant theory, evolution in general relativity (at least in the spatially compact case) is generated by the constraints, thus implying that Dirac observables are constants of motion. Hence finding enough Dirac observables is equivalent to finding a complete set of constants of motion for the Einstein equations, a task which is well nigh impossible. Therefore, to achieve progress in interpreting the formalism, ideas are needed as to how to obtain a physical interpretation of physical states in the absence of explicit Dirac observables.

In contrast to this state of affairs for full blown general relativity, an adequate quantum description of weak gravitational fields *is* available. Sufficiently weak gravitational fields are described by a linearization of general relativity about flat spacetime. Quantization of this (approximate) description of the exact physics is in terms of the graviton Fock representation. Since the exact description is that of full quantum gravity, Fock states in linearised gravity which describe weak gravitational fields must correspond, at least approximately, to states in full quantum gravity. Note that only those Fock states in which the gravitational field has small expectation values and fluctuations describe weak ‘quantum’ gravitational fields

¹ We ignore here, the thorny issues of the applicability of the Copenhagen interpretation to quantum gravity.

and it is only such states which are expected to have full quantum gravity correspondents. An archetypal example of such a state is the graviton vacuum which is expected to provide an adequate physical description of small quantum gravitational fluctuations about flat spacetime.

Let us suppose that within the LQG framework, we find a way to associate *kinematical* structures with Fock states describing weak gravitational fields in the context of a mathematically well defined approximation scheme. Since kinematical structures play a key role in defining physical states, we may hope that the nature of the approximation scheme suggests how to extend this correspondence (of appropriate Fock states with kinematical structures) to physical states. Such a correspondence would provide a way to interpret these physical states (as small quantum perturbations of Minkowski spacetime) in the absence of Dirac observables.

Motivated by these remarks, our aim in this work is to make a preliminary investigation of the relation between quantum linearised gravity and LQG by finding structures in kinematical LQG which correspond to the Fock vacuum. As we shall see, a reasonable definition of such a correspondence requires the satisfaction of infinitely many conditions by the putative LQG structures and therefore the identification of these structures is quite involved.

We proceed as follows. In section 2, we analyse the conditions under which classical linearised gravity is an adequate approximation to full general relativity and show that the approximation may be characterised by a small parameter, ϵ , constructed out of the physically relevant classical distance scales.² Thus, the equations of motion of classical linearised theory are obtained by an expansion of the equations of motion of full general relativity about flat spacetime in which only terms of linear order in ϵ are retained. For the linearised theory vacuum to be of physical relevance, it is necessary that vacuum expectation values and fluctuations of the gravitational field be of order ϵ .³ We analyse this requirement in the context of the r -Fock representation of linearised gravity [1, 2, 3] and show that it leads to a specification of ϵ in terms of an intertwining of the physically relevant

² We are indebted to Abhay Ashtekar for his crucial inputs without which the analysis and results of section 2 would not have been possible.

³ We only expect this condition to hold when the vacuum is probed at ‘large distance scales. A detailed discussion of these matters is given in section 2

classical distance scales with the distance scales relevant to the quantum theory, namely the Planck length and the scale r characterising the r - Fock representation. We use the r - Fock representation because of its structural similarity to the representation used for LQG. The r - Fock representation of linearised gravity is constructed in [1, 3] (see also Appendix 2 of [4]) and its relation to the usual Fock representation is discussed in [3].

In section 3, we propose to identify states in kinematical LQG with the r - Fock vacuum by demanding that the action of the linearised theory operators (to order ϵ) on the r - Fock vacuum be mirrored by the action of their non-linear counterparts in LQG. We show that this demand leads to infinitely many consistency conditions and the consequent necessity of identifying an infinite dimensional set of Fock states with their counterparts in LQG. We also argue that these counterparts are kinematical *distributions* i.e. they are kinematically non-normalizable states which lie in the algebraic dual to the finite span of extended spin network states (for a definition of the latter see [5]). We characterise this identification by a map $\mathcal{M}$ ⁴ from such distributions to appropriate subspaces of r -Fock space. The material after section 3 is devoted to showing the existence of at least one such map which satisfies the consistency conditions.

In section 4 we define two auxilliary structures in LQG which go into our construction of $\mathcal{M}$. The first is a distributional ‘weave’ state which appropriately captures the notion of classical flat space used in the linearised theory. The second is a semi-norm on the space of distributions. The kernel of $\mathcal{M}$ is (partially) specified by those distributions which are of order δ_ϵ in this semi- norm, where δ_ϵ is a small ϵ - dependent parameter. In this section we also specify the general structure of our putative solutions to the consistency conditions. In section 5 we construct such solutions explicitly thus exhibiting distributional states in LQG which correspond to the r - Fock vacuum. In section 6 we argue that the constraints of the linearised theory, represented as operators in the full theory, map the distributional states we constructed in section 5 to the kernel of $\mathcal{M}$ i.e. their images are of order δ_ϵ . Section 8 is devoted to a discussion of our results and a few important identities are proved in the Appendices.

⁴ We actually use a *1 parameter family of maps*, $\mathcal{M}^\epsilon$, in section 3. For pedagogical reasons we gloss over this subtlety and use notation appropriate to a single map $\mathcal{M}$ in this section.

Although our work suggests a strategy to solve the constraints of full gravity order by order, it is pertinent to make a couple of cautionary remarks here to place our results in proper perspective.

(a) Our results would be stronger if we could prove them in the context of a norm rather than a semi- norm. In fact, most of the considerations of sections 5 and 6 *do* go through in the context of a suitably defined norm. The only failure in this context is that of our states to be mapped into the kernel of $\mathcal{M}$ by the linearised Gauss law constraint. In section 7 we comment on this in relation to the fact that the connection variables of linearised theory and LQG take values in distinct Lie algebras (namely $U(1)^3$ and $SU(2)$).

(b) As noted earlier the r - Fock construction is structurally similar to LQG. As briefly discussed in section 7 our constructions of $\mathcal{M}$, the semi- norm as well as the distribution corresponding to the vacuum state, all lean heavily on the existence of similar structures in the r - Fock representation. This is the reason that our constructions ‘work’ at a mathematical level. However, they are still *ad hoc* because they do not arise from any deep *physical* insight.

In view of (a) and (b) above, it is unlikely that the detailed choices of auxilliary structures made in this work lead to a direct relation between “solutions to the constraints to n th order” and exact physical states in nonperturbative LQG. Nevertheless, our hope is that further analysis may lead to a more physically motivated implementation of the general strategy (described in section 3) suggested by our work. If so, this would provide a plausible interpretation of certain states in LQG (namely those corresponding to the vacuum of linearised gravity) without recourse to Dirac observables.

There have been other attempts to identify the vacuum state in LQG [6, 7, 8, 9] as well as to understand how *classical* spacetime emerges from LQG structures [6, 10, 11]. Indeed, these questions are the focus of alot of current research in the field. The related problem of interpretation of physical states in LQG is a very crucial and hard problem. Though the current set of ideas (including ours) may be critiscized on several fronts, we believe that only an exploration of a diversity of such proposals will suggest some way to attack this problem.

Finally, we note that this work is an attempt to implement earlier proposals of Iwasaki and

Rovelli [12].⁵ Indeed, this work would not have been possible without their considerations. It is our view that their beautiful ideas were considerably in advance of the technical state of art of the field and consequently were difficult to implement with precision. We hope that we have partially remedied this by our work here.

In what follows we shall assume familiarity with the loop representation of linearised gravity as discussed in [3]. We set $c = G = 1$ and denote the Planck length by l_P . We also define $O(\epsilon^n)$ and $O^\infty(\epsilon)$ as follows. Let x be an ϵ -dependent complex number. $x = O(\epsilon^n)$ iff $\lim_{\epsilon \rightarrow 0} \frac{|x|}{\epsilon^n}$ exists. $x = O^\infty(\epsilon)$ iff as $\epsilon \rightarrow 0$, $x \rightarrow 0$ faster than any positive power of ϵ .

II. LINEARISED GRAVITY AS AN APPROXIMATE PHYSICAL DESCRIPTION.

A. A quick review of linearised gravity in terms of connections.

The classical Hamiltonian formulation underlying LQG is discussed in [14]. The phase space variables are a spatial $SU(2)$ connection, $A_a^i(\vec{x})$ and a densitized triad field $E_j^b(\vec{y})$. Here a, b denote spatial components, i, j denote internal $SU(2)$ Lie algebra components. To define the linearised theory about a flat background, the spatial topology is chosen to be R^3 . We fix, once and for all, a cartesian coordinate system $\{\vec{x}\}$ on R^3 as well as an orthonormal basis in the Lie algebra of $SU(2)$. Henceforth all components refer to this cartesian coordinate system and to this internal basis. We linearise the $SU(2)$ formulation about the phase space point $(A_a^i = 0, E_i^a = \delta_i^a)$. As in [3], we denote the linearised triad field by e_i^a so that

$$E_i^a = \delta_i^a + e_i^a. \quad (2.1)$$

Since the background connection vanishes, there is no need to introduce a new symbol for the linearised connection. The only non-vanishing Poisson bracket is

$$\{A_a^i(\vec{x}), e_j^b(\vec{y})\}_L = \gamma \delta_a^b \delta_j^i \delta(\vec{x}, \vec{y}). \quad (2.2)$$

⁵ Note that J. Zegwaard independently explored the same issues as Iwasaki and Rovelli using a slightly different approach in his work [13].

and the linearised Gauss Law, Vector and Scalar constraints are

$$G_i^L = \partial_a e_i^a + \epsilon_i^{ja} A_{aj}, \quad (2.3)$$

$$V_a^L = f_{ab}^a, \quad (2.4)$$

$$C^L = \epsilon^{abc} f_{abc}. \quad (2.5)$$

Here γ is the Barbero-Immirzi parameter, $f_{ab}^i = \partial_a A_b^i - \partial_b A_a^i$ is the linearised curvature, ∂_a denotes the flat derivative operator which annihilates the background triad and we use the background triad δ_i^a to freely interchange internal and spatial indices. The linearised $SU(2)$ connection transforms as a triplet of $U(1)$ connections under transformations generated by G_L^i .

Given any oriented piecewise analytic loop β in R^3 with (arbitrary) parametrization s and tangent vector $\dot{\beta}^a$, we define the loop form factor

$$X_\beta^a(\vec{x}) := \oint_\beta ds \delta^3(\vec{\beta}(s), \vec{x}) \dot{\beta}^a, \quad (2.6)$$

and its Fourier transform

$$X_\beta^b(\vec{k}) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int d^3x X_\beta^b(\vec{x}) e^{-i\vec{k} \cdot \vec{x}} = \frac{1}{(2\pi)^{\frac{3}{2}}} \oint_\beta ds e^{-i\vec{k} \cdot \vec{\beta}(s)} \dot{\beta}^a \quad (2.7)$$

For any positive parameter r with dimensions of length, we define the Gaussian smeared loop form factor

$$X_{\beta(r)}^a(\vec{x}) := \int d^3y X_\beta^a(\vec{y}) \frac{e^{-\frac{|\vec{x}-\vec{y}|^2}{2r^2}}}{(2\pi r^2)^{\frac{3}{2}}} \quad (2.8)$$

$$X_{\beta(r)}^a(\vec{k}) = e^{-\frac{k^2 r^2}{2}} X_\beta^a(\vec{k}). \quad (2.9)$$

For any triplet of loops $\alpha_i, i = 1..3$ we define $X_{\alpha}^{ab}(\vec{x})$ and $X_{\alpha(r)}^{ab}(\vec{x})$ via their Fourier transforms to be,

$$X_{\alpha}^{ab}(\vec{k}) := X_{\alpha_i}^a(\vec{k}) \delta^{ib}, \quad X_{\alpha(r)}^{ab}(\vec{k}) := e^{-\frac{k^2 r^2}{2}} X_{\alpha}^{ab}(\vec{k}). \quad (2.10)$$

We also define $G_{\alpha(r)}^{ab}(\vec{x})$ via its Fourier transform,

$$G_{\alpha(r)}^{ab}(\vec{k}) := k X_{\alpha(r)}^+(\vec{k}) (1 + 2i\gamma) m_a m_b - k X_{\alpha(r)}^-(\vec{k}) (1 - 2i\gamma) \bar{m}_a \bar{m}_b. \quad (2.11)$$

Here $m_a, \bar{m}_a$ form the standard transverse basis in momentum space and $X_{\alpha(r)}^\pm$ are the positive and negative helicity components of the transverse, traceless, symmetric part of

$X_{\alpha(r)}^{ab}$. Finally we define the following Dirac observables for the linearised theory:

$$h_{\alpha} = \exp i \int d^3x X_{\alpha}^{ab} A_{ab} = \prod_{k=1}^3 \exp i \oint_{\alpha^k} A_a^k dx^a \quad (2.12)$$

and

$$g_{\alpha(r)} = \int d^3x G_{\alpha(r)}^{ab} e_{(r)ab}, \quad (2.13)$$

where $e_{(r)ab}$ is obtained from e_{ab} by Gaussian smearing. h_{α} and $g_{\alpha(r)}$ play a key role in the considerations of [3] and will continue to play a key role in this work.

B. Classical linearised gravity as an approximate physical description.

Linearised gravity is expected to provide an approximate description of the physics of weak gravitational fields. Such fields are characterised by low curvature. Since curvature is dimensionful, a notion of its smallness requires the introduction of some parameter s_{curv} which has dimensions of length. The restriction to small curvatures implies that $|\partial_a \partial_b e_i^c| < s_{curv}^{-2}$. On dimensional grounds we expect phase space data for low curvatures to satisfy

$$|\partial_a e_i^b|, |A_b^i| < s_{curv}^{-1}. \quad (2.14)$$

Clearly, the larger s_{curv} is, the better is the physical description provided by linearised gravity.

However, small curvatures are only a necessary condition for the application of linearised gravity. There exist nonlinear configurations of the gravitational field, namely large black holes, which have vast regions of small curvature but also possess global features such as horizons. Such global features cannot be described by linearised gravity. More quantitatively, for data satisfying (2.14) we estimate the gravitational energy in a region of size s to be $M = (\frac{1}{s_{curv}})^2 s^3$. If $M \sim s$ the gravitational configuration is expected to be close to that of a black hole. Hence, we require that $M \ll s$ or, equivalently, that $\frac{s}{s_{curv}} \ll 1$. Thus, we expect linearised gravity to be a good description of the gravitational field when s_{curv} is large and when we probe regions of size s much smaller than s_{curv} . For a fixed probe scale s , we define $\epsilon = \frac{s}{s_{curv}}$. Then linearised gravity is a valid description when $\epsilon \ll 1$ and equation (2.14) reads

$$|\partial_a e_i^b|, |A_b^i| < \frac{\epsilon}{s}. \quad (2.15)$$

In addition, we shall assume that the ‘observable’ part of the linearised triad, namely its positive and negative helicity parts, also have spatial variations characterised by the scale s . Specifically, we assume

$$|\partial_c e_{ab}^\pm|, |(-\nabla^2)^{\frac{1}{2}} e_{ab}^\pm| < \frac{\epsilon}{s}. \quad (2.16)$$

In view of their key role in the quantum theory, it is useful to encode the conditions (2.15), (2.16) as conditions on the observables $h_{\boldsymbol{\alpha}}, g_{\boldsymbol{\alpha}(r)}$. From (2.15) we obtain $|\sum_{k=1}^3 \oint_{\alpha^k} A_a^k dx^a| < \frac{\epsilon}{s} |\boldsymbol{\alpha}|$, where $|\boldsymbol{\alpha}|$ is the sum of the lengths of the loops $\alpha^k, k = 1..3$. In order to get a useful bound on $h_{\boldsymbol{\alpha}}$, as well as to ensure that the classical probes (in this case the loops α^i) do not extend into regions of volume much larger than s^3 , we restrict attention to *loops of length at most equal to s* . Then equations (2.12) and (2.15) imply that

$$|h_{\boldsymbol{\alpha}} - 1| < 3\epsilon. \quad (2.17)$$

With the same restriction on loop lengths, we obtain, using (2.7) - (2.11) and (2.16)

$$\begin{aligned} |g_{\boldsymbol{\alpha}(r)}| &= \left| \int X_{\boldsymbol{\alpha}(r)}^{ab}(\vec{x}) [(1 + 2i\gamma)(-\nabla^2)^{\frac{1}{2}} e_{(r)ab}^+(\vec{x}) + (1 - 2i\gamma)(-\nabla^2)^{\frac{1}{2}} e_{(r)ab}^-(\vec{x})] \right| \\ &< 2\sqrt{1 + 4\gamma^2} \epsilon. \end{aligned} \quad (2.18)$$

Note that the above bound is independent of r .

C. Quantum considerations

Equations (2.17) and (2.18) imply that only those states of quantum linearised theory in which the expectation values and fluctuations of the operators $\hat{h}_{\boldsymbol{\alpha}} - 1, \hat{g}_{\boldsymbol{\alpha}(r)}$ are of order ϵ , are physically relevant descriptions of quantum aspects of weak gravitational fields. Since in this work we are interested only in the vacuum state, we require that the vacuum fluctuations and vacuum expectation values of these operators be small (i.e. of $O(\epsilon)$).

As mentioned in the introduction we use the r -Fock representation and the r -Fock vacuum, $|0_r\rangle$ to investigate the consequences of this requirement. We denote the vacuum expectation value of an operator $\hat{O}$ by $\bar{O} := \langle 0_r | \hat{O} | 0_r \rangle$ and its vacuum fluctuation by $\Delta O := \sqrt{\langle 0_r | \hat{O}^\dagger \hat{O} | 0_r \rangle - |\bar{O}|^2}$. From [3], we have that $\bar{g}_{\boldsymbol{\alpha}(r)} = 0$ and that

$$(\Delta g_{\boldsymbol{\alpha}(r)})^2 \leq \frac{1 + 4\gamma^2}{32\pi^3} l_P^2 l_{\boldsymbol{\alpha}_{max}}^2 \int d^3 k k e^{-k^2 r^2} \quad (2.19)$$

$$\leq \frac{1 + 4\gamma^2}{16\pi^2} \frac{l_P^2 s^2}{r^4}, \quad (2.20)$$

where $l_{\alpha_{max}} \leq s$ is the length of the longest loop in the triplet α . To ensure that $\Delta g_{\alpha(r)} = O(\epsilon)$ we set,

$$\frac{l_P s}{r^2} = \epsilon. \quad (2.21)$$

It can be checked that this identification of ϵ ensures that the vacuum expectation values and vacuum fluctuations of $\hat{h}_{\alpha} - 1$ are also at most of $O(\epsilon)$.⁶

How should we interpret equation (2.21)? Note that the parameter r plays no essential role in classical theory. In particular the considerations of section 2.1 are unaltered if we set $r = 0$. However, for quantum operators to be well defined in a (Fock-like representation of) a quantum *field* theory, it is necessary for the basic quantum fields to be appropriately smeared in order to avoid the infinities coming from quantum fluctuations at arbitrarily small distance scales. Here, this smearing is characterised by the distance scale r . While we require that operator fluctuations be small (i.e. of $O(\epsilon)$), the quantum uncertainty principle dictates that operator fluctuations cannot, in general, be arbitrarily small. The potential conflict between these two requirements results in the relation (2.21) between the classically relevant distance scale $s_{curv} = \frac{s}{\epsilon}$ and the distance scales relevant to quantum theory, namely the Planck length l_P and the smearing scale r . Equation (2.21) can be rewritten in the form $r^2 = l_P s_{curv}$. In this form it is clear that if the smearing scale is too small, the resulting short distance quantum fluctuations exceed the upper bounds (2.17) and (2.18). Note that we may choose the individual parameters r , s and s_{curv} to vary with ϵ in different ways. Since we would like to have a non-empty set of probes when $\epsilon \rightarrow 0$, we shall henceforth restrict attention to probe sizes s which do not vanish as $\epsilon \rightarrow 0$. Then equation (2.21) implies that r (and s_{curv}) diverge as $\epsilon \rightarrow 0$. This means that to ensure that the quantum fluctuations in curvature become arbitrarily small, the relevant operators have to be smeared at arbitrarily large distance scales. In this work we shall think of r as a short distance smearing scale relative to the probe scale s and restrict attention to the case $s \gg r$.

⁶ Equation (2.20) only provides an upper bound. In appendix E we construct α^k such that $\Delta g_{\alpha(r)} \approx \epsilon$ to nontrivial leading order in ϵ .

III. A STRATEGY TO IDENTIFY THE VACUUM STATE IN LQG.

In this section we discuss the identification of states in LQG which correspond to the r -Fock vacuum of linearised theory in terms of a map between (suitably defined) states in LQG and (subsets of) r -Fock space. Section 3A contains a few pertinent observations which serve to motivate the definition and desired properties of this map as outlined in section 3B. In section 3C we describe our strategy for an explicit construction of a map which satisfies the criteria of section 3B.

A. A few remarks by way of motivation.

(i) *Linearised theory observables and their LQG counterparts:* The operators $\hat{h}_{\alpha}, \hat{g}_{\alpha(r)}$ play a key role in the identification of the r -Fock vacuum in linearised theory (see [3]). Therefore it is reasonable to look for their counterparts in the $SU(2)$ theory. Using equation (2.15) and equation (2.1), it is straightforward to check that

$$-i(h_{\alpha} - 1) = 2 \sum_{k=1}^3 \text{tr} H_{\alpha^k} \tau^k + O(\epsilon^2), \quad (3.1)$$

$$g_{\alpha(r)} = G_{\alpha(r)}. \quad (3.2)$$

Here H_{α} is the $SU(2)$ holonomy $H_{\alpha} = \frac{1}{2} P \exp - \oint_{\alpha} A_a^i \tau^i$. $\tau^i = -\frac{i}{2} \sigma^i$, where σ^i are the Pauli matrices. Thus τ^i correspond to the fixed basis of the Lie algebra of $SU(2)$ (see section 2) in its defining 2 dimensional representation. Note that (3.1) holds independent of which point of α^k is chosen as the base point. Also,

$$G_{\alpha(r)} := \int d^3x G_{\alpha(r)ab}(\vec{x}) (E_{(r)}^{ab}(\vec{x}) - \delta^{ab}) \quad (3.3)$$

$E_{(r)}^{ab}$ is defined from the $SU(2)$ triad variable by

$$E_{(r)}^{ab}(\vec{x}) := \frac{1}{(2\pi r^2)^{\frac{3}{2}}} \int d^3y e^{-\frac{|\vec{x}-\vec{y}|^2}{2r^2}} E_i^a(\vec{y}) \delta^{bi}. \quad (3.4)$$

We shall refer to the set of functions in linearised theory on the left hand side of equations (3.1) and (3.2) as O_L^{ϵ} and their $SU(2)$ counterparts on the right hand side of these equations as O_F^{ϵ} . As noted at the end of section 2, r is a function of ϵ through (2.21). Since some of observables (namely the ones depending on the triad) depend explicitly on

r , we have a superscript ϵ to signify this dependence. Note also that the linearised theory holonomy operators have an implicit dependence on r coming from the r -dependence of the representation.

Recall that $O_L^\epsilon, O_F^\epsilon$ are labelled by loops of size at most s . Since we are interested in probing regions of size of order s (see section IIB), we further restrict these loops to lie within a volume $(Ss)^3$ about the origin for some positive parameter S *independent of* ϵ .

(ii) *The distributional nature of LQG ‘vacuum’ states:* It is reasonable to expect that any putative vacuum state in the $SU(2)$ theory must be such that the expectation values and fluctuations of the operators $\hat{O}_F^\epsilon$ in this state reproduce the (r - Fock) vacuum expectation values and fluctuations of the operators $\hat{O}_L^\epsilon$ to accuracy ϵ . Since $\hat{O}_F^\epsilon$ are not Dirac observables for full gravity, they are only defined at the (non $SU(2)$ invariant) kinematic level. Note that the $SU(2)$ states cannot lie in the LQG kinematic Hilbert space $\mathcal{L}^2(\bar{\mathcal{A}}, d\mu_0)$ [15, 16] for the following reasons. First, as argued in [17], expectation values and fluctuations for uncountably many $SU(2)$ holonomy operators cannot be reproduced by states in the kinematic Hilbert space and second, as can be easily verified, the smeared triad operator (constructed by replacing E_i^a by $\hat{E}_i^a$ in equation (3.4)) is not well defined on the kinematic Hilbert space. As we shall see later, the $SU(2)$ states lie in (an appropriate subspace) of the algebraic dual, Φ_{kin}^* , to a suitable dense subspace of the kinematic Hilbert space. Since, at least to our knowledge, there is no natural inner product on Φ_{kin}^* , an identification of the vacuum state therein which does not involve explicit evaluation of expectation values and fluctuations of $\hat{O}_F^\epsilon$ is desirable.

(iii) *The approximate nature of the identification between linearised and LQG structures:* Finally we note that since linearised gravity is an approximate description of the exact physics of weak gravitational fields, we expect the correspondence between the r -Fock vacuum and its $SU(2)$ counterparts to be an approximate one.

B. Definition and properties of the map $\mathcal{M}^\epsilon$.

Motivated by the above remarks, we define the identification of suitable $SU(2)$ distributions with the r - Fock vacuum, $|0_r\rangle$, as follows. Recall that equation (2.21) implies that r is a function of ϵ . Hence $|0_r\rangle$ corresponds to a 1 parameter family of states, 1 state for

each $r(\epsilon)$, $0 < \epsilon \ll 1$. Consider a corresponding 1 parameter family of putative $SU(2)$ vacuum states Ψ_0^ϵ . We require the existence of a 1 parameter family of maps, $\mathcal{M}^\epsilon$, which map the 1 parameter family of states Ψ_0^ϵ to the 1 parameter family of *sets of* r - Fock states $\{|0_r > + |\phi^\epsilon >, \sqrt{\langle \phi^\epsilon | \phi^\epsilon \rangle} = O(\epsilon^2)\}$. We denote this requirement as

$$\mathcal{M}^\epsilon \Psi_0^\epsilon = |0_r > + O(\epsilon^2). \quad (3.5)$$

Thus, for each fixed ϵ , $\mathcal{M}^\epsilon$ is a map from a subspace of Φ_{kin}^* to (a subset of) the set of all subsets of r - Fock space. Using similar notation, we require that Ψ_0^ϵ be such that the action of any $\hat{O}_F^\epsilon$ on Ψ_0^ϵ mirrors that of the corresponding $\hat{O}_L^\epsilon$ on $|0_r >$:

$$\mathcal{M}^\epsilon \hat{O}_F^\epsilon \Psi_0^\epsilon = \hat{O}_L^\epsilon |0_r > + O(\epsilon^2). \quad (3.6)$$

Since we can always measure a linear combination of the basic linearised operators $\hat{O}_L^\epsilon$, we impose the following ‘linearity’ condition on $\mathcal{M}^\epsilon$. Let $\hat{O}_{iL}^\epsilon, \hat{O}_{iF}^\epsilon, i = 1..N$ be a set of N linearised and ‘full’ operators and let a_0, a_i be a set of $N + 1$ complex parameters each of $O(1)$ with N itself being independent of ϵ . Then we require that

$$\mathcal{M}^\epsilon \left(\sum_{i=1}^N a_i \hat{O}_{iF}^\epsilon + a_0 \right) \Psi_0^\epsilon = \left(\sum_{i=1}^N a_i \hat{O}_{iL}^\epsilon + a_0 \right) |0_r > + O(\epsilon^2). \quad (3.7)$$

Although the above equation does not involve an explicit determination of $\bar{O}_F^\epsilon, \Delta O_F^\epsilon$, it is, nevertheless, motivated by the requirement that $\bar{O}_L^\epsilon, \Delta O_L^\epsilon$ be approximated by their $SU(2)$ counterparts. To see this, note that if we could identify the states Ψ_0^ϵ , the above equation serves to define the ‘minimal’ subspace $\mathcal{S}^{\epsilon*} \subset \Phi_{kin}^*$ which can serve as the domain of the map $\mathcal{M}^\epsilon$ i.e.

$\mathcal{S}^{\epsilon*} = \{\Psi^\epsilon \in \Phi_{kin}^*, \Psi^\epsilon = (\sum_{i=1}^N a_i \hat{O}_{iF}^\epsilon + a_0) \Psi_0^\epsilon\}$. Let us suppose that there exists an inner product on this space which is well approximated by the r -Fock inner product $\langle, \rangle$ in the following sense. Consider $\Psi_1, \Psi_2 \in \mathcal{S}^{\epsilon*}$ with $\mathcal{M}^\epsilon \Psi_j = |\psi_j > + O(\epsilon^2), j = 1, 2$. Then let $\langle, \rangle_F$ on $\mathcal{S}^{\epsilon*}$ be such that

$$\langle \Psi_1, \Psi_2 \rangle_F = \langle \psi_1 | \psi_2 \rangle + O(\epsilon^2)(\sqrt{\langle \psi_1 | \psi_1 \rangle} + \sqrt{\langle \psi_2 | \psi_2 \rangle}) + O(\epsilon^4). \quad (3.8)$$

It is straightforward to check that for such a $\langle, \rangle_F$, the equations (3.5)-(3.7) imply that fluctuations and expectation values of $\hat{O}_F^\epsilon$ in any state Ψ_0^ϵ approximate their linearised counterparts to the desired accuracy of $O(\epsilon)$.

The existence of ambiguities of $O(\epsilon^2)$ in the right hand sides of (3.5) and (3.6) may be motivated as follows. Equation (3.6) may be thought of as the quantum version of the classical relations (3.1) and (3.2). Since $\hat{O}_L^\epsilon |0_r\rangle$ has norm of $O(\epsilon)$, the ambiguity in the image of $\mathcal{M}^\epsilon$ must be of higher order than ϵ and we choose it to be $O(\epsilon^2)$.

The rest of this work is devoted to showing the existence of at least one (1 parameter family of maps) $\mathcal{M}^\epsilon$ which satisfies equations (3.5)-(3.7).⁷

C. Strategy for an explicit construction of $\mathcal{M}^\epsilon$

The approximate correspondence between linearised and full gravity motivated us to require that the image of $\mathcal{M}^\epsilon$ be defined with small ambiguities of $O(\epsilon^2)$ in equations (3.5)-(3.7). Similar considerations should also apply to the domain of $\mathcal{M}^\epsilon$ i.e. it seems plausible that $\mathcal{M}^\epsilon$ should map 2 distributions which differ by a ‘small’ amount to the same image. Our ignorance of any useful inner product on Φ_{kin}^* makes it difficult to define this notion of ‘smallness’. However, even though we do not know of any useful inner product we can define norms and semi- norms on Φ_{kin}^* . Indeed, we shall define a semi- norm on Φ_{kin}^* and (partially) specify the ‘kernel’ of $\mathcal{M}^\epsilon$ as containing distributions which are of $O(\delta_\epsilon)$ in this semi- norm for a suitably defined small ϵ dependent parameter, δ_ϵ . We define the kernel of (the 1 parameter family of maps) $\mathcal{M}^\epsilon$ to be the set (of all 1 parameter families) of distributions, whose images by $\mathcal{M}^\epsilon$ are 1 parameter families of sets of states with r - Fock norm of $O(\epsilon^2)$.

The operators $\hat{O}_L^\epsilon$ satisfy the following relation in their actions on $|0_r\rangle$ (this relation is just the ‘Poincar’e invariance condition’ of [3] evaluated to $O(\epsilon)$):

$$i(\hat{h}\boldsymbol{\alpha} - 1)|0_r\rangle = \hat{g}_{\boldsymbol{\alpha}(r)}|0_r\rangle + O(\epsilon^2). \quad (3.9)$$

This in conjunction with (3.6) implies that

$$\mathcal{M}^\epsilon(2(\sum_{k=1}^3 tr \hat{H}_{\alpha^k} \tau_k) + \hat{G}_{\boldsymbol{\alpha}(r)})\Psi_0^\epsilon = O(\epsilon^2). \quad (3.10)$$

⁷ Strictly speaking, our results are slightly weaker in that we analyse these equations subject to a mild restriction on the loops labelling $O_L^\epsilon, O_F^\epsilon$, namely that these loops intersect the weave defined in section 4A in at most a finite number of points.

Our strategy will be to attempt to solve the following ‘ $SU(2)$ Poincaré invariance condition’ upto ambiguities of $O(\delta_\epsilon)$ ⁸:

$$2 \sum_{k=1}^3 \text{tr} \hat{H}_{\alpha^k} \tau_k \Psi_0^\epsilon = -\hat{G}_{\alpha(r)} \Psi_0^\epsilon + O(\delta_\epsilon), \quad (3.11)$$

to obtain Ψ_0^ϵ . Once we obtain such a Ψ_0^ϵ , we can define the action of $\mathcal{M}^\epsilon$ as follows:

$$\mathcal{M}^\epsilon \Psi_0^\epsilon := |0_r\rangle + O(\epsilon^2) \quad (3.12)$$

$$\mathcal{M}^\epsilon \hat{G}_{\alpha(r)} \Psi_0^\epsilon := \hat{g}_{\alpha(r)} |0_r\rangle + O(\epsilon^2). \quad (3.13)$$

Equation (3.11) ensures that the relevant conditions involving the connection dependent operators in the linearised and $SU(2)$ theories are satisfied. Finally, $\mathcal{M}^\epsilon$ can be extended by appropriate linearity to satisfy equation (3.7).

IV. THE WEAVE AND THE SEMI- NORM.

The semi- norm comprises one of two auxilliary structures that we will define in order to construct an $\mathcal{M}^\epsilon$ satisfying equations (3.5)- (3.7). It turns out that in order to define the semi- norm we use, we need to define a second auxilliary structure namely a (distributional) ‘weave’ state. This state is the counterpart of the flat triad δ_i^a of linearised theory. We define the weave state in section 4A and the semi- norm in section 4B. In section 4C, we describe the general structure of the $SU(2)$ distributions which we shall consider in the rest of this work.

A. The weave state.

The weave state is the analog of the classical flat triad δ_i^a in the following sense. We shall construct a set of extended spin network states denoted by $\{|\psi_\Delta\rangle\}$ which are all based on the same ‘lattice-like’ graph Δ . Then the weave state,⁹ $\Psi_{weave} \in \Phi_{kin}^*$, has the following

⁸ Actually, we solve an equation of the form $((2 \sum_{k=1}^3 \text{tr} \hat{H}_{\alpha^k} \tau_k) + \hat{G}_{\alpha(r)} + a_\alpha^\epsilon) \Psi_0^\epsilon = O(\delta_\epsilon)$ where a_α^ϵ is complex and of $O(\epsilon^2)$. Note that $a_\alpha^\epsilon \Psi_0^\epsilon$ is also in the kernel of $\mathcal{M}^\epsilon$.

⁹ In referring to this state as a weave state, we follow the nomenclature used for states which correspond to classical spatial geometries [10]. The state defined here is based on a graph identical to that used by Rovelli in [4] and Zegwaard in [13].

property

$$(\hat{E}_{i(r)}^a(\vec{x})\Psi_{weave})[|\psi_\Delta\rangle] = (1 + O^\infty(\epsilon))\delta_i^a\Psi_{weave}[|\psi_\Delta\rangle]. \quad (4.1)$$

Here, $\vec{x}$ is restricted to be within the probe region i.e. $|\vec{x}| < Ss$ (see 3A(i) for a definition of S) and $O^\infty(\epsilon)$ has been defined at the end of section 1. Moreover for any state $|\psi_{\Delta_\perp}\rangle$ orthogonal to the span of $\{|\psi_\Delta\rangle\}$, the weave state satisfies

$$(\hat{E}_{i(r)}^a(\vec{x})\Psi_{weave})[|\psi_{\Delta_\perp}\rangle] = \delta_i^a\Psi_{weave}[|\psi_{\Delta_\perp}\rangle] = 0. \quad (4.2)$$

In subsection 1 below, we describe our construction of the weave state and in subsection 2, we display some of its properties. Calculations pertaining to the explicit verification of (4.1) and (4.2) are contained in Appendix A.

1. Construction of the weave state

Let Δ be the graph corresponding to a cubical lattice with edges along the the cartesian coordinate directions. Let the lattice spacing be $\sqrt{\frac{\gamma l_P^2}{2}}$ and let it occupy a volume $(2L)^3$ centered about the origin. Let L be such that $L \gg s$ and $s_{curv} \gg L \gg r \gg l_P$.¹⁰

Define the spin net $|\Delta\rangle$ based on the graph Δ by:

- (a) Coloring all edges of Δ with $j = \frac{1}{2}$ i.e. with spin-half representations.
- (b) Defining trivial intertwiners at the vertices of Δ i.e. mapping, trivially, the $j = \frac{1}{2}$ representation on an incoming edge in the i th direction ($i = 1, 2, 3$) at any vertex to the $j = \frac{1}{2}$ representation on the outgoing edge in the same direction (this pertains to vertices which are not at the boundary of Δ).
- (c) Choosing *any* intertwiners at the vertices on the boundary of the lattice i.e. ‘tying up’ the boundary points on the planes $x^i = \pm L$ in any convenient way. This choice has no bearing on our subsequent considerations.

¹⁰ Note that since these inequalities are ϵ dependent, Δ and Ψ_{weave} also depend on ϵ . However, we shall regard this dependence as ‘weak’ in the sense that we hope that a better treatment incorporating constructions appropriate to asymptotic flatness (see for example [18]) will allow L to be taken to ∞ without altering our conclusions. In this connection, we note that the inequality $s_{curv} \gg L$ is not crucial to the considerations of appendix A; as can be verified, $L \gg s_{curv}$ would suffice equally well. In what follows, we shall refrain from explicitly denoting the $L(\epsilon)$ dependence of Ψ_{weave}

Define the spin net $|\Delta, \{\vec{x}_k, k = 1..N\} >$ as follows. Pick N points $\{\vec{x}_k, k = 1..N\}$ on the graph Δ such that they are *not* on the vertices of Δ . In addition to (a), (b) and (c) above, define the intertwiners at the new vertices $\{\vec{x}_k, k = 1..N\}$ as follows. Let $\vec{x}_k$ be on an edge along the i th direction. Then the intertwiner is chosen as $2\tau_{iA}^B$, $A, B = 1, 2$ (clearly τ_{iA}^B maps the incoming $j = \frac{1}{2}$ representation to the outgoing $j = \frac{1}{2}$ representation at the vertex $\vec{x}_k$). Note that in terms of the labelling of extended spin nets defined in [5], $\tau_{iA}^B = M_a^i C_A^{aB}$ where (i) C_A^{aB} is the invariant tensor which maps the $j = \frac{1}{2}$ representation to the product of the $j = \frac{1}{2}$ and $j = 1$ representations and (ii) M_a^i , $i = 1, 2, 3$ are vectors in the $j = 1$ representation namely, $M_a^1 = (\sqrt{\frac{3}{8}}i, 0, -\sqrt{\frac{3}{8}}i)$, $M_a^2 = (-\sqrt{\frac{3}{8}}, 0, \sqrt{\frac{3}{8}}i)$ and $M_a^3 = (0, \sqrt{\frac{3}{2}}i, 0)$. $A, B = 1, 2$ are $j = \frac{1}{2}$ indices and $a = 1, 2, 3$ is a $j = 1$ index.

It can be checked that $|\Delta, \{\vec{x}_k, k = 1..N\} >$ is normalised and that

$$< \Delta, \{\vec{y}_j, j = 1..M\} | \Delta, \{\vec{x}_k, k = 1..N\} > = 0, \quad (4.3)$$

unless $\{\vec{y}_j\} = \{\vec{x}_k\}$. Ψ_{weave} is defined as the distribution

$$\Psi_{weave} := \sum_{N=0}^{\infty} \sum_{\vec{x}_k, k=1..N} (-i)^N < \Delta, \{\vec{x}_k\} |, \quad (4.4)$$

where the second (uncountable) sum is over all possible values of $\{\vec{x}_k\}$. Clearly, Ψ_{weave} is in Φ_{kin}^* i.e. it has a well defined action on any finite linear combination of spinnets.

2. Properties of the weave state.

The weave state has nice properties with respect to the action of the smeared triad operators. The considerations of Appendix A show that equation (4.1) is satisfied for $|\vec{x}| < Ss$ (see 3A(i) for a definition of S). Although we do not display them here, straightforward calculations along the lines of Appendix A show that for $|\vec{x}_j| < Ss, j = 1..n$, n independent of ϵ ,

$$\begin{aligned} \prod_{j=0}^{n-1} \hat{E}_{(r)i_{n-j}}^{a_{n-j}}(\vec{x}_{n-j}) \Psi_{weave}[\psi_{\Delta} >] &= \hat{E}_{(r)i_n}^{a_n}(\vec{x}_n) \dots \hat{E}_{(r)i_1}^{a_1}(\vec{x}_1) \Psi_{weave}[\psi_{\Delta} >] \\ &= (1 + O^{\infty}(\epsilon)) \prod_{j=0}^{n-1} \delta_{i_{n-j}}^{a_{n-j}} \Psi_{weave}[\psi_{\Delta} >]. \end{aligned} \quad (4.5)$$

Here $|\psi_\Delta \rangle = |\Delta \rangle$ or $|\Delta\{\vec{x}_k\} \rangle$. From (4.5) it is straightforward to show that

$$\prod_{j=0}^{n-1} (\hat{E}_{(r)i_{n-j}}^{a_{n-j}}(\vec{x}_{n-j}) - \delta_{i_{n-j}}^{a_{n-j}}) \Psi_{weave}[|\psi_\Delta \rangle] = O^\infty(\epsilon) \Psi_{weave}[|\psi_\Delta \rangle]. \quad (4.6)$$

Equation (4.6) captures the precise sense in which the weave state is the analog of the flat triad δ_i^a *within the probe region* $|\vec{x}| < Ss$. Note however that since $G_{\alpha_{ab}(r)}$ is not, in general, a function of compact support (see Appendix B), the Poincare invariance condition (3.9) as well as its $SU(2)$ counterpart (3.11) involve properties of the triad everywhere *including* $|\vec{x}| > Ss$. Hence to proceed further either (a) we need the weave state to capture properties of the flat triad for all $\vec{x}$ or (b) we may attempt to restrict the Poincare invariance condition in such a way as to depend only on properties of the triad in the probe region. For simplicity we shall assume that option (a) is viable. More precisely, we assume that an improved treatment will yield Ψ_{weave} which satisfies (4.6) for all $\vec{x}$ (see also footnote 10) in this regard). While this seems to us to be a reasonable assumption, we emphasize that its validity must be checked in a more comprehensive treatment. We also note that preliminary calculations made by us indicate that option (b) may also be viable. Since the relevant analysis pertaining to option (b) is a bit involved and may obscure our main line of argument in an already involved paper, we refrain from reporting on this matter here.

Thus, we shall assume that (4.6) is valid everywhere on the spatial slice. Then, as shown in appendix B the following identities hold for n independent of ϵ :

$$\prod_{j=0}^{n-1} \hat{G}_{\alpha_{n-j}}^\dagger \Psi_{weave}[|\Delta \rangle] = O^\infty(\epsilon) \Psi_{weave}[|\Delta \rangle], \quad (4.7)$$

$$\prod_{j=0}^{n-1} \hat{G}_{\alpha_{n-j}}^\dagger \Psi_{weave}[|\Delta\{\vec{x}_k\} \rangle] = O^\infty(\epsilon) \Psi_{weave}[|\Delta\{\vec{x}_k\} \rangle]. \quad (4.8)$$

B. The semi- norm.

As discussed in [3], r - Fock states can be thought of as $U(1)^3$ distributions i.e. distributions with respect to the finite span of $U(1)^3$ flux network states. This is in close structural similarity to the LQG states under consideration, the latter being $SU(2)$ distributions with respect to the finite span of $SU(2)$ spinnets. Although we do not show it here, our choice of

semi- norm below is motivated by the existence of a similar structure in the $U(1)^3$ theory. We shall comment on this further in section 7. For now, we shall simply display our choice of semi- norm and show in the subsequent section that our choices furnish the map $\mathcal{M}^\epsilon$ in accord with the strategy outlined in section 3C.

Consider the set of graphs G_Δ which intersect Δ at most in a finite number of points. Denote the spinnet based on the trivial graph as $|\cdot\rangle$. Clearly, every spinnet based on a graph $\alpha \in G_\Delta$ can be written in the form $\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}}|\cdot\rangle$ where $\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}}$ is the generalised holonomy operator corresponding to the spin net labelled by the graph α , invariant tensors $\vec{c}$ and representation vectors $\vec{M}$ [5] (the labels defining the colour of the edges of α have been suppressed as in [5]). Explicitly, in the connection representation, $|\cdot\rangle$ corresponds to the state $\psi(\bar{A}) = 1$ and the action of the operator $\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}}$ corresponds to multiplication by $\mathcal{N}_{\alpha, \vec{c}, \vec{M}}(\bar{A})$ (see equation (4.19) of [5]).

For any such $\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}}$ consider the states $\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}}|\Delta\rangle$ and $\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}}|\Delta\{\vec{x}_k, k = 1..N\}\rangle$. Denote by V the finite span of these states for all choices of extended spinnets based on every $\alpha \in G_\Delta$. Denote by $V_\perp$ the set of states which are orthogonal to every element of V . Clearly $V + V_\perp$ is a dense subspace of the kinematical Hilbert space.

Next, consider the set of *loops* $G_{\Delta loop}^s$ which are of length less than or equal to s ¹¹ and which intersect Δ at most in a finite number of points. For $\eta_I, \beta_J \in G_{\Delta loop}^s$, $I = 1..P$, $J = 1..M$, $k_J \in \{1, 2, 3\}$, define the operators

$$\hat{O}_{\{\eta_I, \beta_J, k_J\}} := \prod_{I=1}^P \frac{\text{Tr} \hat{H}_{\eta_I}}{2} \prod_{J=1}^M \text{Tr} \hat{H}_{\beta_J} \tau_{k_J} \quad \hat{O}_{\{\eta_I\}} := \prod_{I=1}^P \frac{\text{Tr} \hat{H}_{\eta_I}}{2}. \quad (4.9)$$

Define $V_{loop}^s \subset V$ to be the finite span of the states

$\{\hat{O}_{\{\eta_I, \beta_J, k_J\}}|\Delta\rangle, \hat{O}_{\{\eta_I\}}|\Delta\rangle, \hat{O}_{\{\eta_I, \beta_J, k_J\}}|\Delta\{\vec{x}_k, k = 1..N\}\rangle, \hat{O}_{\{\eta_I\}}|\Delta\{\vec{x}_k, k = 1..N\}\rangle\}$ for all choices of $\{\eta_I, \beta_J, k_J, \vec{x}_k\}$ and finite P, M, N .

For any $\Psi \in \Phi_{kin}^*$ we define the semi- norm

$$||\Psi||^\epsilon = \sup_{|\psi\rangle} |\Psi[|\psi\rangle]| \quad (4.10)$$

where $|\psi\rangle$ ranges over the following:

(a) $|\psi\rangle$ is a spinnet in $V_\perp$,

¹¹ Note that unlike the loops which label $O_L^\epsilon, O_F^\epsilon$, these loops are *not* restricted to lie within a volume $(Ss)^3$ about the origin (see section 3A(i)).

(b) $|\psi\rangle$ is $(M + P)^{-(M+P)} \hat{O}_{\{\eta_I, \beta_J, k_J\}} |\Delta\rangle$ or $P^{-P} \hat{O}_{\{\eta_I\}} |\Delta\rangle$ or $(M + P)^{-(M+P)} \hat{O}_{\{\eta_I, \beta_J, k_J\}} |\Delta\{\vec{x}_k, k = 1..N\}\rangle$ or $P^{-P} \hat{O}_{\{\eta_I\}} |\Delta\{\vec{x}_k, k = 1..N\}\rangle$, for all choices of $\{\eta_I, \beta_J, k_J, \vec{x}_k\}$ and finite P, M, N .

$\|\cdot\|^\epsilon$ is a semi-norm precisely because $|\psi\rangle$ is not allowed to range over the orthogonal complement of V_{loop}^s in V . The above semi-norm is ϵ -dependent both due to the (weak) dependence of Δ on ϵ (see footnote 10) as well as the dependence of $s \gg r$ on ϵ . The ϵ -dependence of the semi-norm is denoted by the superscript ϵ .

C. General structure of the LQG distributions under study.

Consider an arbitrary distribution $\Phi \in \Phi_{kin}^*$. Define $\Psi \in \Phi_{kin}^*$ from Φ, Ψ_{weave} as follows.

- (a) Let $|\psi\rangle \in V_\perp$ be a spinnet. Then $\Psi[\psi] := 0$.
- (b) Let $\alpha \in G_\Delta$ and consider any ‘spinnet’ operator $\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}}$ as defined in section 4.2 above. Then $\Psi[\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}} |\Delta\rangle := \Psi_{weave}[|\Delta\rangle \Phi[\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}} \cdot\rangle]$ and $\Psi[\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}} |\Delta\{\vec{x}_k\}\rangle := \Psi_{weave}[|\Delta\{\vec{x}_k\}\rangle \Phi[\hat{\mathcal{N}}_{\alpha, \vec{c}, \vec{M}} \cdot\rangle]$. This defines the action of Ψ on a basis of V .
- (c) Extend the action of Ψ as defined in (a) and (b) to the finite span of all spinnets by linearity. The $SU(2)$ states of interest in the rest of this work will have the structure of Ψ defined above.

V. THE VACUUM AS AN $SU(2)$ DISTRIBUTION.

Below, we exhibit solutions to the $SU(2)$ Poincare invariance condition (3.11) (modulo terms in the kernel of $\mathcal{M}^\epsilon$ (see footnote 8, section 3C in this regard)), thus implementing the strategy of section 3C. The particular choice of our solutions as well as that of δ_ϵ is motivated by their analogs in the description of r -Fock states by $U(1)^3$ distributions. We shall comment on this further in section 7. Here we shall simply display our choices and show that they furnish the required $\mathcal{M}^\epsilon$.

A. Explicit form of the solution to the Poincare invariance condition.

The putative solution Ψ_0^ϵ is of the type described in section 4C with $\Phi = \Phi_0^\epsilon$ where Φ_0^ϵ is defined as

$$\begin{aligned}\Phi_0^\epsilon &= \delta_{A=0} + \hat{E}^2 \delta_{A=0}, \\ \hat{E}^2 &:= \frac{i}{2\gamma l_P^2} \int d^3 k k [(1 + 2i\gamma)(\hat{E}_{(r)}^+(\vec{k}))^\dagger \hat{E}_{(r)}^+(\vec{k}) - (1 - 2i\gamma)(\hat{E}_{(r)}^-(\vec{k}))^\dagger \hat{E}_{(r)}^-(\vec{k})],\end{aligned}\quad (5.1)$$

where $\delta_{A=0}$ is the distribution corresponding to the connection $A_a^i = 0$ i.e. for any $\phi(\bar{A}) \in \mathcal{L}^2(\bar{\mathcal{A}}, d\mu_0)$,

$$\delta_{A=0}[\phi(\bar{A})] := \phi(\bar{A})|_{\bar{A}=0}. \quad (5.2)$$

Consider a set of operators $\hat{O}_i, i = 0..n$ such that each $\hat{O}_i$ is of the type defined in equation (4.9). Then we have the following key identity which is proved in Appendix D.

$$\hat{O}_0 \hat{G} \alpha_{1(r)} \hat{O}_1 \dots \hat{G} \alpha_{n(r)} \hat{O}_n \Psi_0^\epsilon = \Psi^\epsilon + O^\infty(\epsilon). \quad (5.3)$$

Here Ψ^ϵ is a distribution of the type defined in section 4C with $\Phi = \Phi^\epsilon$ where Φ^ϵ is defined as $\Phi^\epsilon := \hat{O}_0(\int G \alpha_{1(r) a_1 b_1} \hat{E}_{(r)}^{a_1 b_1}) \hat{O}_1 \dots (\int G \alpha_{n(r) a_n b_n} \hat{E}_{(r)}^{a_n b_n}) \hat{O}_n \Phi_0^\epsilon$.

The above identity (5.3) states that distribution on its left hand side is the sum of Ψ^ϵ and a distribution of semi-norm $O^\infty(\epsilon)$. Since Ψ^ϵ is also of the type defined in section 4C, this identity allows us to rewrite the Poincare invariance condition as a condition on the ‘ Φ ’ part of the distribution. In this regard, it is useful to define the following seminorm, $|| \cdot ||_1^\epsilon$, pertinent to Φ :

$$||\Phi||_1^\epsilon = \sup_{|\phi\rangle} |\Phi[|\phi\rangle]|, \quad (5.4)$$

where $|\phi\rangle \in \{(M+P)^{-(M+P)} \hat{O}_{\{\eta_I, \beta_J, k_J\}} |\cdot\rangle, P^{-P} \hat{O}_{\{\eta_I\}} |\cdot\rangle\}$ for all choices of $\{\eta_I, \beta_J, k_J\}$ and finite P, M, N ($\hat{O}_{\{\eta_I, \beta_J, k_J\}}, \hat{O}_{\{\eta_I\}}$ are defined by (4.9)). Note that for any distribution Ψ of the type defined in section 4C, we have that

$$||\Psi||^\epsilon = ||\Phi||_1^\epsilon. \quad (5.5)$$

Then it follows from (5.3) that the $SU(2)$ Poincare invariance condition (3.11) on Ψ_0^ϵ , for loops $\alpha^k \in G_{\Delta loop}^s$ (and which are, as in (3.11) confined to a volume $(Ss)^3$ about the origin), is equivalent to the following condition on Φ_0^ϵ :

$$2 \sum_{k=1}^3 \text{tr} \hat{H}_{\alpha^k} \tau_k \Phi_0^\epsilon = - \left(\int d^3 x G \alpha_{(r) ab}(\vec{x}) \hat{E}_{(r)}^{ab}(\vec{x}) \right) \Phi_0^\epsilon + O(\delta_\epsilon). \quad (5.6)$$

Here the last term denotes a distribution with seminorm $|| ||_1^\epsilon$ of $O(\delta_\epsilon)$. In the next subsection we show that Φ_0^ϵ does indeed satisfy (5.6) upto a term of $O(\epsilon^2)$ which is in the kernel of $\mathcal{M}^\epsilon$. More precisely, we show that

$$2 \sum_{k=1}^3 \text{tr} \hat{H}_{\alpha^k} \tau_k \Phi_0^\epsilon = - \left(\int d^3x G_{\alpha(r)ab}(\vec{x}) \hat{E}_{(r)}^{ab}(\vec{x}) \right) \Phi_0^\epsilon + \epsilon^2 c_{\alpha} \Phi_0^\epsilon + O(\delta_\epsilon), \quad (5.7)$$

where c_{α} is a complex number of $O(1)$ depending only on α . Since $\mathcal{M}^\epsilon \epsilon^2 c_{\alpha} \Phi_0^\epsilon = \epsilon^2 c_{\alpha} |0_r\rangle$, this extra term is in the kernel of $\mathcal{M}^\epsilon$. This is equivalent to showing that Ψ_0^ϵ satisfies (3.10) for $\text{tr} \hat{H}_{\alpha^k} \tau_k, \hat{G}_{\alpha(r)}$ labelled by loops α^k each of which are of length at most equal to s , are confined to a volume $(Ss)^3$ about the origin, *and* which intersect Δ in at most a finite number of points. Note that this is a slightly weaker condition than (3.10) which was defined for all loops with length at most s confined to a volume $(Ss)^3$ about the origin *irrespective of their intersections with Δ* . We shall only impose this slightly weaker condition of Poincare invariance on Ψ_0^ϵ rather than the stronger one (3.10).

B. Verification of (5.7).

We set $\delta_\epsilon = \epsilon^4$. We shall explain this choice later in this subsection. From (C5) it follows that

$$\hat{E}_{(r)i}^a(\vec{k}) \left(2 \sum_{k=1}^3 \text{tr}(H_{\alpha^k}(A) \tau^k) \right) |_{A=0} = -i l_P^2 \gamma X_{(r)\alpha^i}^a(\vec{k}). \quad (5.8)$$

Also, setting $n = 0, N = 1$ and choosing $O_1^H = \text{tr}(H_{\alpha^k}(A) \tau^k)$ in (C3) it follows that

$$\hat{E}^2 \sum_{k=1}^3 \text{tr}(H_{\alpha^k}(A) \tau^k) |_{A=0} = \epsilon^2 c_{\alpha}, \quad (5.9)$$

where c_{α} is a constant of $O(1)$ which depends only on the triplet of loops α .

The action of the left hand side of (5.7) on $|\phi\rangle$ of the type appearing in (5.4) is

$$L.H.S. := \sum_{k=1}^3 \text{tr}(H_{\alpha^k}(A) \tau^k) \phi(A) |_{A=0} + \hat{E}^2 \sum_{k=1}^3 \text{tr}(H_{\alpha^k}(A) \tau^k) \phi(A) |_{A=0} \quad (5.10)$$

where we have denoted the ket $|\phi\rangle$ by the corresponding wave function in the connection representation, $\phi(A)$. Since $\text{tr}(H_{\alpha^k}(A) \tau^k)$ vanishes at $A = 0$, the first term in (5.10) vanishes and in the second term either both triad operators of $\hat{E}^2$ act on $\text{tr}(H_{\alpha^k}(A) \tau^k)$ or one acts

on $tr(H_{\alpha^k}(A)\tau^k)$ and one on $\phi(A)$. Using (5.8) and (5.9) to evaluate these contributions, we obtain

$$\begin{aligned}
L.H.S. &= -(\int G\alpha_{(r)ab}\hat{E}_{(r)}^{ab})^\dagger\phi(A)|_{A=0} + \epsilon^2 c\alpha\phi(A)|_{A=0} \\
&= -(\int G\alpha_{(r)ab}\hat{E}_{(r)}^{ab})^\dagger\phi(A)|_{A=0} + \epsilon^2 c\alpha(\Phi_0^\epsilon[\phi] - \hat{E}^2\delta_{A=0}[\phi]) \\
&= -(\int G\alpha_{(r)ab}\hat{E}_{(r)}^{ab})^\dagger\phi(A)|_{A=0} + \epsilon^2 c\alpha\Phi_0^\epsilon[\phi] + O(\epsilon^4),
\end{aligned} \tag{5.11}$$

where we have used (C3) to obtain the $O(\epsilon^4)$ term. Using (C3) again, the action of the right hand side of (5.7) on ϕ is

$$R.H.S. = -(\int G\alpha_{(r)ab}\hat{E}_{(r)}^{ab})^\dagger\phi(A)|_{A=0} + O(\epsilon^4). \tag{5.12}$$

This completes the verification of (5.7).

Note that from (C2) and (C3) it is easy to see that $(\int d^3x G\alpha_{(r)ab}(\vec{x})\hat{E}_{(r)}^{ab}(\vec{x}))\Phi_0^\epsilon$ is $O(\epsilon^2)$. But the image of this quantity by $\mathcal{M}^\epsilon$ is expected to be (see equation(3.13)) $\hat{g}\alpha_{(r)}|0_r >$. Since the norm of the latter is expected to be of $O(\epsilon)$ from (2.20), we do not want $(\int d^3x G\alpha_{(r)ab}(\vec{x})\hat{E}_{(r)}^{ab}(\vec{x}))\Phi_0^\epsilon$ to be in the kernel of $\mathcal{M}^\epsilon$. Hence we choose $\delta_\epsilon \ll \epsilon^2$ and our choice is $\delta_\epsilon = \epsilon^4$.

This argument is merely a plausability one. Strictly speaking, we need to show that for every α^k such that $\Delta g\alpha_{(r)} \approx \epsilon$ to nontrivial leading order in ϵ , there exists $|\phi >$ of the type in (5.4) such that $(\int d^3x G\alpha_{(r)ab}(\vec{x})\hat{E}_{(r)}^{ab}(\vec{x}))\Phi_0^\epsilon[|\phi >] \approx \epsilon^2$ to nontrivial leading order in ϵ . It is straightforward to see that this is indeed true for $|\phi > = \sum_{k=1}^3 tr(\hat{H}_{\alpha^k}\tau^k)|\cdot >$. This suggests that we generalise the definition of the seminorm so as to allow $|\phi >$ of this type. While we do not anticipate any obstruction to such a trivial generalisation, we leave a thorough verification of this to future work.

C. Higher order operators.

It can be verified that the action of quadratic and higher order products of operators of type $\hat{O}_L^\epsilon$ on $|0_r >$ yields states of norm at most $O(\epsilon^2)$ (this is under the assumption that the number of operators of type $\hat{O}_L^\epsilon$ in the product is independent of ϵ). Therefore, it is natural to demand that the action of their $SU(2)$ counterparts map the $SU(2)$ vacuum into the kernel of $\mathcal{M}^\epsilon$. Also note that the $SU(2)$ function, $(Tr(H_\alpha) - 1)$, $\alpha \in G_{\Delta loop}^s$ has

leading order term quadratic in the connection. Hence, it is natural to also demand that $((Tr\hat{H}_\alpha) - 1)\Psi_0^\epsilon$ be in the kernel of $\mathcal{M}^\epsilon$.

Indeed, using the key identities (5.3), (C2) and (C3), it is straightforward to verify that (i) $((Tr\hat{H}_\alpha) - 1)\Psi_0^\epsilon = \epsilon^2 d_\alpha \Psi_0^\epsilon + O(\delta_\epsilon)$, where d_α is an α -dependent complex number of $O(1)$ and $\alpha \in G_{\Delta loop}^s$.

(ii) The action of any product of 2 operators each of which belongs to the set $\{O_F^\epsilon, ((Tr\hat{H}_\alpha) - 1), \alpha \in G_{\Delta loop}^s\}$, on Ψ_0^ϵ , yields distributions in the kernel of $\mathcal{M}^\epsilon$. Specifically, these distributions are either of type $O(\epsilon^2)\Psi_0^\epsilon$ or of $O(\delta_\epsilon)$.

(iii) The action of any product of n operators ($n > 2$, n independent of ϵ) each of which is in the set defined in (ii) above, on Ψ_0^ϵ , yields a distribution of $O(\delta_\epsilon)$.

VI. THE LINEARISED CONSTRAINTS IN THE $SU(2)$ THEORY.

We present qualitative arguments to show that the $SU(2)$ vacuum state is mapped to a distribution of $O(\delta_\epsilon)$ by the linearised constraints (2.3)-(2.5) expressed as operators in the $SU(2)$ theory.

We first discuss the purely connection dependent linearised vector and scalar constraints V^L and C^L . We shall impose these constraints everywhere on the spatial slice except on the set of measure zero containing points on the weave Δ . This implies that in the $SU(2)$ theory it should be possible to regularise $\hat{V}^L, \hat{C}^L$ in terms of holonomies along edges which do not intersect Δ . From the structure of Ψ_0^ϵ defined in section 4C, this implies that the action of the regularised operators (we denote these by $\hat{V}_{reg}^L, \hat{C}_{reg}^L$) on Ψ_0^ϵ is simply to change Φ_0^ϵ to $\hat{V}_{reg}^L \Phi_0^\epsilon, \hat{C}_{reg}^L \Phi_0^\epsilon$. Note also that classically, the positive and negative helicity parts of the triad Poisson commute with V^L, C^L . Hence in any reasonable regularization, we expect that in the limit of the regulator being removed, we have that

$$[\hat{V}^L, \hat{E}^2] = [\hat{V}^L, \hat{E}^2] = 0. \quad (6.1)$$

Finally, since V^L, C^L vanish when evaluated at the zero connection, we expect that

$$\hat{V}^L \delta_{A=0} = \hat{C}^L \delta_{A=0} = 0. \quad (6.2)$$

Equations (6.1) and (6.2) imply that $\hat{V}^L, \hat{C}^L$ annihilate Φ_0^ϵ , and in view of the above discussion also annihilate Ψ_0^ϵ .

Next, we examine the linearised Gauss Law constraint G_i^L (see (2.3). Since in our work the basic triad operator is $\hat{E}_{(r)i}^a$, we define the Gaussian smeared constraint $G_{(r)i}^L$ by

$$G_{(r)i}^L(\vec{x}) = \int d^3y G_i^L(\vec{y}) \frac{e^{-\frac{|\vec{x}-\vec{y}|^2}{2r^2}}}{(2\pi r^2)^{\frac{3}{2}}} \quad (6.3)$$

$$= \partial_a e_{(r)i}^a(\vec{x}) + \epsilon_i^{ja} A_{(r)aj}(\vec{x}). \quad (6.4)$$

The first term in (6.4) is obtained after a by parts integration and $A_{(r)aj}(\vec{x})$ is the Gaussian smeared connection. In the $SU(2)$ theory our arguments pertaining to V^L, C^L apply to the connection dependent part of (6.4). Hence we expect that for any reasonable definition of the connection part of $G_{(r)i}^L(\vec{x})$, this connection part annihilates Ψ_0^ϵ . Hence we need analyse only the triad dependent part of (6.4).

At the classical level we would like to impose the linearised constraints as $O(\epsilon)$ restrictions on the phase space. Since (6.3) has the dimensions of inverse length, we need to integrate it against a gauge parameter $\Lambda^i(\vec{x})$ which has dimensions of inverse length square so as to obtain the dimensionless quantity $G_{(r)}^L(\Lambda) := \int d^3x \Lambda^i G_{(r)i}^L$. We denote the triad dependent part of $G_{(r)}^L(\Lambda)$ by

$$G_{(r)e}^L(\Lambda) := - \int d^3x (\partial_a \Lambda^i(\vec{x})) e_{(r)i}^a(\vec{x}), \quad (6.5)$$

where we have performed a by parts integration and assumed that $\Lambda^i(\vec{x})$ is of compact support. We shall meet the requirement $|G_{(r)}^L(\Lambda)| \sim O(\epsilon)$ by imposing the (sufficient) condition

$$\int d^3x |\partial_a \Lambda^i(\vec{x})| \sim O(1). \quad (6.6)$$

In the $SU(2)$ theory we have

$$\hat{G}_{(r)e}^L(\Lambda) \Psi_0^\epsilon = - \int d^3x (\partial_a \Lambda^i(\vec{x})) (\hat{E}_{(r)i}^a(\vec{x}) - \delta_i^a) \Psi_0^\epsilon. \quad (6.7)$$

Using (4.1) and (6.6), we obtain

$$|\hat{G}_{(r)e}^L(\Lambda) \Psi_{weave}[|\psi_\Delta >]| = O^\infty(\epsilon), \quad (6.8)$$

where $|\psi_\Delta >| \in \{|\Delta >, |\Delta\{\vec{x}_k\} >|\}$. From equation (6.8) and the structure of Ψ_0^ϵ as defined in section 4C, we have that

$$\hat{G}_{(r)e}^L(\Lambda) \Psi_0^\epsilon = O(\delta_\epsilon) \Leftrightarrow \hat{G}_{(r)E}^L(\Lambda) \Phi_0^\epsilon = O(\delta_\epsilon) \quad (6.9)$$

where $\hat{G}_{(r)E}^L(\Lambda) := - \int d^3x (\partial_a \Lambda^i(\vec{x})) \hat{E}_{(r)i}^a(\vec{x})$ and where $O(\delta_\epsilon)$ in the first equality is defined with respect to $|| \cdot ||^\epsilon$ and in the second equality with respect to $|| \cdot ||_1^\epsilon$ (see (5.4)).

Using the (inverse Fourier transform of) equation (C5) it is straightforward to obtain

$$\hat{G}_{(r)E}^L(\Lambda) \text{tr}(H_\alpha(A) \tau^i)|_{A=0} = - \int (\partial_a \Lambda^i) X_{(r)\alpha}^a = \int (\partial_a X_{(r)\alpha}^a) \Lambda^i = 0, \quad (6.10)$$

where we have used the fact that the smeared loop form factor is divergence free i.e.

$$\partial_a X_{(r)\alpha}^a = 0. \quad (6.11)$$

Moreover, equation (C5) also implies that

$$\hat{G}_{(r)E}^L(\Lambda) \text{tr}(H_\alpha(A))|_{A=0} = 0. \quad (6.12)$$

Equations (6.10) and (6.12) imply that

$$\delta_{A=0}[\hat{G}_{(r)E}^L(\Lambda)|\phi >] = 0 \quad (6.13)$$

where $|\phi >$ is of the type defined below (5.4).

Using methods similar to those in Appendix C, it is straightforward to verify that for $O_\alpha^H \in \{\text{tr}(H_\alpha \tau^i), \frac{\text{tr} H_\alpha}{2}, \alpha \in G_{\Delta loop}^s\}$ we have that,

$$|\hat{E}^2 \hat{G}_{(r)E}^L(\Lambda) O_\alpha^H(A)|_{A=0} < \epsilon^2 d_0 l_P^2 \gamma \int d^3x |\partial_a \Lambda_i(\vec{x})| |X_{(r)\alpha}^a(\vec{x})|, \quad (6.14)$$

where d_0 is defined in (C3). Using (6.6) and the bound

$$|X_{(r)\alpha}^a(\vec{x})| = |\oint_\alpha ds \dot{\alpha}^a e^{\frac{-|\vec{x}-\vec{\alpha}(s)|^2}{2r^2}}| < \frac{s}{(2\pi r^2)^{\frac{3}{2}}}, \text{ we obtain}$$

$$|\hat{E}^2 \hat{G}_{(r)E}^L(\Lambda) O_\alpha^H(A)|_{A=0} < \epsilon^2 d_0 \frac{s l_P^2 \gamma}{r^3} = O(\epsilon^4) \frac{r}{s} \quad (6.15)$$

From the above equation and the fact that $s \gg r$ it follows straightforwardly that

$$|\hat{E}^2 \hat{G}_{(r)E}^L(\Lambda) \prod_{I=1}^N O_I^H(A)|_{A=0} < N^3 O(\epsilon^4). \quad (6.16)$$

Finally, it is easily verified that the second equality in equation (6.9) follows from (6.13) and (6.16), thus implying that the first equality in (6.9) holds. This completes our arguments in support of the hypothesis that the linearised constraints expressed as operators in the $SU(2)$ theory map the $SU(2)$ vacuum to the kernel of $\mathcal{M}^\epsilon$.

VII. GENERALIZATION TO A NORM?

In this section, we enquire as to whether our considerations admit a generalization to a norm. We find that the linearised Gauss constraint provides the only obstruction to the most obvious such generalization. In order to obtain insight into why this happens, we are led to a comparison between the $SU(2)$ structures hitherto defined and the $U(1)^3$ structures (alluded to in section 1) which provided the motivation for these definitions.

$||^\epsilon$ is not a norm because the states $|\psi\rangle$ defined in (a) and (b) of section 4B do not span the kinematic Hilbert space. This can be remedied by requiring that the loops η_I, β_J in section 4B (a), (b) be replaced by *edges*. More precisely, define $G_{\Delta edge}^s$ such that every element of $G_{\Delta edge}^s$ is an analytic edge of length at most s which intersects Δ in at most a finite number of points. Since higher spin representations can be built up as products of spin $\frac{1}{2}$ representations and since edge holonomies along longer edges are products of those along shorter edges, it follows that $|\psi\rangle$ in section 4B (a), (b) defined for $\eta_I, \beta_J \in G_{\Delta edge}^s$ span the kinematic Hilbert space. Thus, with this modification $||^\epsilon$ becomes a norm. Let us denote this norm by $||^\epsilon_{norm}$. Remarkably, all our results with the single exception of the treatment of G_i^L go through in this norm. Specifically, (6.11) no longer holds if α is an edge and not a loop.

As mentioned in section 1, the reason our $SU(2)$ constructions work is due to their structural analogy with the corresponding $U(1)^3$ ones in the description [3] of r -Fock states as $U(1)^3$ distributions. Hence, it is instructive to view the above failure to impose the linearised Gauss law as an approximate constraint in the $SU(2)$ theory in the context of this structural analogy. Therefore we briefly describe the relevant $U(1)^3$ structures below.

(1) From [3], $|0_r\rangle$ may be written as the $U(1)^3$ distribution Φ_0^L defined by

$$\Phi_0^L := \sum_{\alpha, \{\mathbf{q}\}} c_0 \alpha_{\{\mathbf{q}\}} < \alpha, \{\mathbf{q}\} |, \quad (7.1)$$

$$c_0 \alpha_{\{\mathbf{q}\}} = \exp \left(-i \frac{\gamma}{2} \int d^3x G_{\alpha_{\{\mathbf{q}\}}(\mathbf{r})\mathbf{ab}}(\vec{x})^* X_{\alpha_{\{\mathbf{q}\}}(\mathbf{r})}^{ab}(\vec{x}) \right), \quad (7.2)$$

where $|\alpha, \{\mathbf{q}\}\rangle$ is a $U(1)^3$ gauge invariant flux network state based on the triplet of graphs α labelled by the integers $\{\mathbf{q}\}$ and $G_{\alpha_{\{\mathbf{q}\}}(\mathbf{r})\mathbf{ab}}$ is essentially the same as $G_{\alpha, (r)ab}$ defined by

(2.11) in this paper. See [3] for the precise definitions of $G_{\alpha, \{q\}(r)ab}, X_{\alpha, \{q\}(r)}^{ab}$. Note that γ_0 in [3] is related to γ of this work by $\gamma_0 = 2\gamma$.

(2) Consider the algebraic dual Φ_{kin}^{*L} to the finite span of extended (i.e. not necessarily gauge invariant) $U(1)^3$ flux nets. For any $\Phi \in \Phi_{kin}^{*L}$ define the seminorm $||\Phi||_{seminorm}^L = \sup_{|\phi\rangle} |\Phi[|\phi\rangle|]$ with $|\phi\rangle := \hat{h}_{\alpha}|\cdot\rangle_L$, the length of each α^i being at most s , and with $|\cdot\rangle_L$ denoting the flux network state corresponding to the trivial loop. Then the right hand side of (7.1) can be expanded in this semi- norm as

$$\Phi_0^L = \Phi_{0approx}^L + O(\epsilon^4) \quad (7.3)$$

$$\Phi_{0approx}^L := \delta_{A=0} + \hat{e}^2 \delta_{A=0}, \quad (7.4)$$

$$\hat{e}^2 := \frac{i}{2\gamma l_P^2} \int d^3k k [(1 + 2i\gamma)(\hat{e}_{(r)}^+(\vec{k}))^\dagger \hat{e}_{(r)}^+(\vec{k}) - (1 - 2i\gamma)(\hat{e}_{(r)}^-(\vec{k}))^\dagger \hat{e}_{(r)}^-(\vec{k})], \quad (7.5)$$

where $\delta_{A=0}$ is the $U(1)^3$ distribution corresponding to the $U(1)^3$ connection $A_a^i = 0$.

(3) For any $\Phi \in \Phi_{kin}^{*L}$, define the (stronger than $||\cdot||_{seminorm}^L$) seminorm

$$||\Phi||^L := \sup_{|\phi\rangle} |\Phi[|\phi\rangle|]| \quad (7.6)$$

with $|\phi\rangle := \prod_{I=1}^N \frac{\hat{h}_{\alpha_I}|\cdot\rangle_L}{N^{3N}}$ for $\alpha_I^k, k = 1..3, I = 1..N$ being loops of length at most s , N finite. Then $\Phi_{0approx}^L$ satisfies the Poincare invariance condition (3.9) modulo distributions of $O(\epsilon^4)$ in the seminorm (7.6) and modulo terms of type $O(\epsilon^2)\Phi_{0approx}^L$.

From (1)- (3) it is clear as to how $U(1)^3$ structures motivated our definitions of $SU(2)$ structures. In particular the $U(1)^3$ structures $\hat{e}_i^a = -il_P^2 \gamma \frac{\delta}{\delta A_a^i}, \Phi_0^L$ and $||\cdot||^L$ bear a close analogy to the $SU(2)$ structures $\hat{E}_i^a = -il_P^2 \gamma \frac{\delta}{\delta A_a^i}, \Phi_0^\epsilon$ and $||\cdot||^\epsilon$. In this context the failure of $||\hat{G}_{(r)}^L(\Lambda)\Psi_0^\epsilon||^\epsilon$ to be of $O(\delta_\epsilon)$ may be traced, as indicated below, to the difference between the gauge groups $U(1)^3$ and $SU(2)$.

Consider the algebraic dual $\Phi_{kin}^{*L inv}$ to the finite span of gauge invariant $U(1)^3$ fluxnets. Since larger loop $U(1)^3$ holonomies can be constructed as products of smaller loop ones, it follows that $|\phi\rangle$ of the type in (7.6) span the space of $U(1)^3$ gauge invariant fluxnets. Hence, although $||\cdot||^L$ is a seminorm on Φ_{kin}^{*L} , it is a *norm* on $\Phi_{kin}^{*L inv}$. Moreover, since $\hat{g}_{\alpha(r)}, \hat{h}_{\alpha}$ are $U(1)^3$ gauge invariant, the entire treatment of the $U(1)^3$ theory can be done at the gauge invariant level, and hence, in the context of a norm.

In contrast the very notion of linearization of the full $SU(2)$ theory as implemented in [1, 3] involves an $SU(2)$ gauge *variant* background. Since the $SU(2)$ counterparts of $\hat{g}_{\alpha(r)}, \hat{h}_{\alpha}$ are

themselves *not* $SU(2)$ gauge invariant, it seems appropriate to use gauge variant kinematical states $|\psi\rangle$ in the definition of $||\cdot||^\epsilon$. Ultimately, however, only $SU(2)$ *invariant* distributions are physically relevant in LQG. Hence one may further restrict attention in (5.4) to only those $|\phi\rangle$ which are gauge invariant in the hope that the resulting seminorm on Φ_{kin}^* reduces to a norm on Φ_{kin}^{*inv} , where Φ_{kin}^{*inv} is the algebraic dual to the finite span of spin nets which are gauge invariant with respect to $SU(2)$ transformations within the probe region. However, this hope is not realised because, in contrast to the $U(1)^3$ case, due to the non-abelian nature of $SU(2)$, traces of large loop holonomies are not expressible as (sums of) products of traces of small loop holonomies. Moreover, with this restriction, it seems that $||\hat{G}\alpha_{(r)}\Psi_0^\epsilon||^\epsilon = O(\epsilon^4)$ which implies that a change in our choice of δ_ϵ to $\delta_\epsilon \ll \epsilon^4$ would also be needed.

An alternate strategy would be to, in analogy with $U(1)^3$ theory, restrict attention to only such $|\psi\rangle$ in (4.10) which satisfy $\hat{G}_{(r)}^L(\Lambda)|\psi\rangle = 0$ i.e. try to impose the $U(1)^3$ constraint as an $SU(2)$ operator constraint. Our intuition is that no solutions to such an equation exist because $SU(2)$ spinnets are “attuned” to $SU(2)$ gauge transformations rather than $U(1)^3$ ones. Instead, since in the linearised theory we are interested in connections close to $A_a^i = 0$, we may attempt to solve $\hat{G}_{(r)E}^L(\Lambda)\phi(A)|_{A=0} = 0$. Indeed, viewed in this way, $|\phi\rangle$ defining $||\cdot||_1^\epsilon$ are precisely of this type.

On account of the above discussion, our view is that

- (a) the failure of (6.11) in the context of $||\cdot||_{norm}^\epsilon$ is not an accident; rather it is indicative of the difference between the gauge groups $U(1)^3$ and $SU(2)$
- (b) the seminorm $||\cdot||^\epsilon$ seems to be a reasonable structure to use.¹²

Nevertheless, a (putative) perturbative treatment of the constraints would be more powerful in the context of a stronger seminorm. In view of the discussion above it would be interesting to see if we could enlarge the definition of $|\phi\rangle$ in (5.4) such that (i) $\hat{G}_{(r)E}^L(\Lambda)\phi(A)|_{A=0} = 0$ (ii) $|\phi\rangle$ although not themselves necessarily gauge invariant, do span the space of $SU(2)$ gauge invariant spinnets based on graphs in G_Δ .

¹² It may be verified that $||G_{(r)E}^L(\Lambda)\Psi_0^\epsilon||_{norm}^\epsilon = O(\epsilon^2)\frac{r}{s}$. This suggests the possibility of exploiting the freedom in (b) of Appendix A to either redefine $\delta_\epsilon = \epsilon^2\frac{r}{s}$ for $\frac{r}{s} > \epsilon^2$ or to choose $\frac{r}{s} \leq \epsilon^2$. However, since the $O(\epsilon^2)\frac{r}{s}$ bound depends crucially on (in our view the unduly weak premise) equation (6.6), we believe that such a strategy may be inappropriate.

VIII. DISCUSSION

One of the features of our work is the, at first perhaps surprising but nevertheless essential, role of the probe scale s . At the classical level s acts as an infrared cut off, allowing us to ignore large distance effects leading to the formation of black holes. The consequent restriction of the quantum observables $\hat{h}_{\alpha}, \hat{g}_{\alpha(r)}$ to loops α^k of length at most s leads to the determination of ϵ via equation (2.20). Moreover, the derivations of various bounds in this work lead us to believe that (i) it is only for loops with length close to s that the fluctuations of the linearised observables can be close to ϵ ¹³ (ii) the seminorms of the various distributions encountered are determined primarily by the action of these distributions on states of type (b) in section 4B with η_I, β_J of length $\approx s$ and $M, P \approx 1$. Thus, s seems to play a key role in characterising the nature and content of the approximate descriptions at both the classical and the quantum levels.

A second feature of our work is (for reasons described in section 3A (ii)) the use of distributions as opposed to kinematically normalizable states. Since we believe that distributional states will play a key role in future developments, it would be profitable to explore structures on Φ_{kin}^* . In the theory of infinite dimensional vector spaces, topologies are typically defined via (families of) seminorms [19]. In analogy with this and in view of the fact that the seminorm $|| ||^\epsilon$ on Φ_{kin}^* has played a key role in our considerations, we advocate further study of seminorms on the space Φ_{kin}^* .

The seminorm $|| ||^\epsilon$ provided us with the beginnings of a perturbative treatment of the constraints in terms of the expansion parameter ϵ . The first step was to verify that the linearised constraints mapped our states into the kernel of $\mathcal{M}^\epsilon$. In showing this we did not display an explicit regularization of the constraints. Rather, we predicated our arguments on certain expectations of any “reasonable” regularization procedure. The discussion in section 6 indicates that any such reasonable regularization procedure must provide some preferred role to the weave Δ . This seems to be in contrast to the standard Thiemann regularization [20] of say, the full Hamiltonian constraint, in which there is no preferred background structure.

¹³ More work is necessary to establish exactly which α^k yield $\Delta g_{\alpha(r)} \approx \epsilon$ to nontrivial leading order in ϵ . However the bound (2.19) clearly indicates that any such α^k should have lengths $\approx s$.

We believe that this possible conflict between the (putative) perturbative treatment of the constraints and the available nonperturbative treatment is of a generality which transcends the details of our particular constructions. Further work is needed to verify if such a conflict is indeed present in our work and if so, to seek a resolution. Another important open issue is to understand better the relation between the $U(1)^3$ gauge transformations in linearised theory and the $SU(2)$ ones of LQG. The discussion in section 7 was indicative of our lack of understanding of this relation. We feel that a first exploratory step towards more clarity would be to see if we can use a more sophisticated seminorm in accordance with (i)-(ii) at the end of section 7. In this regard, it would be worthwhile to base our discussion of $|\phi\rangle$ in (5.4) in terms of graph-based spinnets rather than loop based wave functions.

The key issue underlying the above paragraph is that of the relation between (putative) perturbative solutions to the constraints and the known structure of the nonperturbative ones as spatially diffeomorphism invariant, $SU(2)$ gauge invariant distributions [21]. As mentioned earlier in this work, we have merely shown the existence of a map $\mathcal{M}^\epsilon$ satisfying the various requirements of section 3. While this was a nontrivial and necessary exercise, it is important to emphasize that our constructions were based primarily on *mathematical* analogies between $U(1)^3$ and $SU(2)$ structures. Without additional physical insight into the significance of the various choices we have made, we are unsure if these choices (or suitable modifications thereof) ensure that (putative) perturbative solutions converge to nonperturbative ones (or, indeed, if they ensure even the existence of a well defined perturbative scheme). Since the issue is a deep and quite general one, clearly more work and thought is needed for further progress. Indeed, the main virtue of our work may be that it offers a concrete and detailed context wherein this issue may be analysed.

It would be of interest to see if our considerations can be generalised to construct LQG correspondents of states other than the linearised theory vacuum. It would also be interesting to adapt our general framework so as to find LQG correspondents of other known exactly solvable restrictions of full gravity such as Einstein- Rosen waves [22]. From equation (3.8) it is tempting to speculate that such ‘consistency’ with a multitude of exactly solvable restrictions of full gravity may lead to information about the scalar product of full quantum gravity.

There have been other efforts to construct the vacuum state in LQG. We comment briefly on three of these, namely [9], [8] and [7]. The recent work [9] is a very interesting attempt to construct the vacuum state as a kinematically normalizable state in LQG rather than, as in this work, a genuine distribution. Due to our unfamiliarity with the detailed considerations of [9], we are unable to comment further on its relation to our work. The work [8] contains a nice description of the author’s beautiful complexifier construction of coherent states. In the LQG context this construction yields genuinely distributional states. However, the author’s viewpoint is that since such states do not seem to support the action of flux operators [8], it is appropriate to use kinematically normalizable ‘cut off’ versions of these states which are similar to the gauge theory coherent states of [6]. As mentioned earlier, our viewpoint is that distributions *are* the appropriate structures. It would be interesting to see if the complexifier generated distributions satisfy the requirements of section 3B in the context of some suitably defined $\mathcal{M}^\epsilon$. Distributional states are also used in [7]. Our work seems to be closest in spirit to [7] and it may be of interest to draw parallels between the graphs chosen as ‘probes’ in [7] and the states in section 4B (b) which go into the definition of the seminorm $|| \cdot ||^\epsilon$.

In closing, we reiterate that many of the considerations of this work were motivated by the earlier work of Iwasaki and Rovelli [12]. Since the work of Zegwaard [13] also tried to address the same problem as [12], it would be worthwhile to revisit it in the light of our work here.

Acknowledgements: We thank Carlo Rovelli and Thomas Thiemann for encouragement. We are indebted to Daniel Sudarsky and Alejandro Perez for their criticisms. We thank Abhay Ashtekar for encouragement and crucial insights as well as Chris Van Den Broeck for numerous discussions. We are grateful for the hospitality of the Center for Gravitational Physics and Geometry, PSU, where we discussed a preliminary version of this work. A significant part of this work was completed when the author was visiting the Ludwig Maximilian University in Munich. We thank Professor Mukhanov of LMU for his kind support during this period. We thank Supurna Sinha for pointing us to the Poisson Sum Formula. We are indebted to Alok Laddha for his comments and for a careful perusal of a draft version of this work.

APPENDIX A

We sketch the main steps involved in showing (4.1). Let ψ_α be any spinnet based on a graph α with edges $e_I, I = 1..N$ (we shall use the notation $|\psi_\alpha\rangle$ interchangeably with ψ_α). Then (see for example equation (3.6) of [5]) we have that

$$\begin{aligned} (\hat{E}_{(r)i}^a(\vec{x}) \Psi_{weave})[\psi_\alpha] &= \Psi_{weave}[\hat{E}_{(r)i}^a(\vec{x})\psi_\alpha] \\ &= i l_P^2 \gamma \sum_I \int dt \dot{e}_I^a \frac{e^{-\frac{|\vec{x}-\vec{e}_I(t)|^2}{2r^2}}}{(2\pi r^2)^{\frac{3}{2}}} \Psi_{weave}[(h_{e_I}(1,t)\tau_i h_{e_I}(t,0))_A^B \frac{\partial \psi_\alpha}{\partial h_{e_I}(1,0)_A^B}] \end{aligned} \quad (A1)$$

Here $h_{e_I}(v, u)$ is the holonomy along the I th edge from parameter value u to parameter value v , the beginning and end points of e_I being at parameter value 0 and 1. Note that the argument of Ψ_{weave} above is obtained from ψ_α by inserting the intertwiner τ_i at the point $\vec{e}_I(t)$ on the I th edge of α . Since $\Psi_{weave} \in \Phi_{kin}^*$, we have that $\Psi_{weave}[(h_{e_I}(1,t)\tau_i h_{e_I}(t,0))_A^B \frac{\partial \psi_\alpha}{\partial h_{e_I}(1,0)_A^B}]$ is finite. Hence we can neglect contributions to the integral in (A1) coming from any sets of measure zero in parameter space. In particular, we can drop contributions from pre-existing vertices $\vec{e}_I(0), \vec{e}_I(1)$.

Lemma 1: If ψ_α is orthogonal to all $|\Delta, \{\vec{x}_k\}\rangle$, then $(\hat{E}_{(r)i}^a(\vec{x})\Psi_{weave})[\psi_\alpha] = 0$.

Proof: Note that ψ_α is orthogonal to $|\Delta, \{\vec{x}_k\}\rangle$ iff any of the following are true:

- (i) α is different from Δ ,
- (ii) $\alpha = \Delta$ but has at least one edge with $j \neq \frac{1}{2}$,
- (iii) $\alpha = \Delta$, every edge is labelled by $j = \frac{1}{2}$ and there exists at least one point in Δ at which the intertwiners for ψ_α and $|\Delta, \{\vec{x}_k\}\rangle$ are orthogonal i.e. $\sum_{i_1..i_t, j_1..j_s} C_{(\alpha)j_1..j_s}^{i_1..i_t} C_{(\Delta, \{\vec{x}_k\})j_1..j_s}^{i_1..i_t} = 0$. Here $C_{(\alpha)}$ is the intertwiner for the vertex in question for ψ_α , $C_{(\Delta, \{\vec{x}_k\})}$ is the intertwiner for the same vertex for $|\Delta, \{\vec{x}_k\}\rangle$ and $i_1..i_t$ label the representations on the incoming edges (for which the vertex is a ‘target’) and $j_1..j_s$ label the representations on the outgoing edges (for which the vertex is a ‘source’). Note that these intertwiners may be further split into a product of invariant intertwiners and representation vectors but this is not necessary for our purposes.

The result (iii) is obtained straightforwardly from the definition of the intertwiners and the orthogonality properties of the representations of $SU(2)$ with respect to the Haar measure. Clearly if any of (i)-(iii) are true of ψ_α , the same is true of $(h_{e_I}(1,t)\tau_i h_{e_I}(t,0))_A^B \frac{\partial \psi_\alpha}{\partial h_{e_I}(1,0)_A^B}$.

except at most on sets of measure zero in parameter space. Thus, for ψ_α orthogonal to all $|\Delta, \{\vec{x}_k\}\rangle$, we have that

$$(\hat{E}_{(r)i}^a(\vec{x})\Psi_{weave})[\psi_\alpha] = 0 = \delta_i^a \Psi_{weave}[\psi_\alpha]. \quad (\text{A2})$$

Lemma 2: Let the extent L of Δ be such that $s_{curv} \gg L \gg r \gg l_P$. Further, let $L \gg s$ and let $\vec{x}$ lie in the probe region of size $(Ss)^3$ about the origin (see 3A(i) for the definition of S) so that $L \gg |\vec{x}|$. Then

$$(\hat{E}_{i(r)}^a(\vec{x})\Psi_{weave})[|\Delta, \{\vec{x}_k\}\rangle] = (1 + O^\infty(\delta) + O^\infty(\tau))\delta_i^a \Psi_{weave}[|\Delta, \{\vec{x}_k\}\rangle] \quad (\text{A3})$$

with $\delta = \frac{\sqrt{l_P^2 \gamma}}{r}$ and $\tau = \frac{r}{L}$.

Proof: From (A1) and the definition of Ψ_{weave} , we have that $(\hat{E}_{i(r)}^a(\vec{x})\Psi_{weave})[|\Delta, \{\vec{x}_k\}\rangle] = 0$ if $a \neq i$. Hence the result must be proportional to δ_i^a . Let us suppose that $a = i = 1$ (our considerations below trivially extend to any other choice of $a = i$). Then from equations (A1) and (4.4), it is straightforward to show that

$$\begin{aligned} \Psi_{weave}[\hat{E}_{i(r)}^a(\vec{x})|\Delta, \{\vec{x}_k\}\rangle] &= (-i)^N \left(\sum_{L_z} \sqrt{\frac{l_P^2 \gamma}{2}} \frac{e^{-\frac{(z-L_z)^2}{2r^2}}}{\sqrt{2\pi r}} \right) \\ &\quad \left(\sum_{L_y} \sqrt{\frac{l_P^2 \gamma}{2}} \frac{e^{-\frac{(y-L_y)^2}{2r^2}}}{\sqrt{2\pi r}} \right) \int_{-L}^L d\vec{x} \frac{e^{-\frac{(x-\vec{x})^2}{2r^2}}}{\sqrt{2\pi r}}. \end{aligned} \quad (\text{A4})$$

Here (L_y, L_z) range over the lattice points in the $x = 0$ plane. Note that from (2.21), the condition $s_{curv} \gg L$ is the same as $r^2 \gg l_P L$. Straightforward estimates using the latter condition and the condition that $L \gg |\vec{x}|$, in conjunction with the Poisson Summation formula¹⁴ [23] yield

$$\left(\sum_{L_z} \sqrt{\frac{l_P^2 \gamma}{2}} \frac{e^{-\frac{(z-L_z)^2}{2r^2}}}{\sqrt{2\pi r}} \right), \left(\sum_{L_y} \sqrt{\frac{l_P^2 \gamma}{2}} \frac{e^{-\frac{(y-L_y)^2}{2r^2}}}{\sqrt{2\pi r}} \right) = 1 + O^\infty(\delta) + O^\infty(\tau). \quad (\text{A5})$$

Also it is easy to check that

$$\int_{-L}^L d\vec{x} \frac{e^{-\frac{(x-\vec{x})^2}{2r^2}}}{\sqrt{2\pi r}} = 1 + O^\infty(\tau). \quad (\text{A6})$$

¹⁴ We are indebted to Supurna Sinha for pointing us to the Poisson Sum formula which plays this key role as an estimation technique in our work.

This completes the proof of Lemma 2.

In order to obtain equation (4.1) from Lemma 2, it suffices to choose $\delta = \epsilon^\alpha$, $\tau = \epsilon^\beta$, $\alpha, \beta > 0$. Below we display 2 explicit choices of ϵ dependence of the various scales involved which satisfy all our requirements.

(a) Choose s independent of ϵ , $r = \epsilon^{-\frac{1}{2}}$, $s_{curv} = \epsilon^{-1}s$, $L = \epsilon^{-\frac{3}{4}}s$. Thus $\alpha = \frac{1}{2}$ and $\beta = \frac{1}{4}$ and we have that $s_{curv} \gg L \gg r \gg s \sim O(1)$.

(b) s dependent on ϵ , $s \neq 0$ as $\epsilon \rightarrow 0$. Choose $\alpha > \frac{1}{2}$ and $\beta > 0$, $\alpha > \beta > \alpha - 1$. Let $s = \epsilon^{1-2\alpha}l_P$, $r = \epsilon^{-\alpha}l_P$, $s_{curv} = \epsilon^{-2\alpha}l_P$, $L = \epsilon^{-(\alpha+\beta)}l_P$. Then for $\alpha > 1$, we have that $s_{curv} \gg L \gg s \gg r$ and for $\alpha < 1$, $s_{curv} \gg L \gg r \gg s$.

As mentioned at the end of section 2, we shall set $s \gg r$ i.e. we shall restrict attention to the choice (b) with $\alpha > 1$.

APPENDIX B

In this appendix we prove the identities (4.7),(4.8) under the assumption that (4.6) holds for all $\vec{x}$ on the spatial slice. From (2.11) we have that

$$G_{\alpha(r)ab}(\vec{x}) = (1 + 2i\gamma)g_{\alpha(r)ab}^{(+)}(\vec{x}) - (1 - 2i\gamma)g_{\alpha(r)ab}^{(-)}(\vec{x}) \quad (\text{B1})$$

where $g_{\alpha(r)ab}^{\pm}$ are the positive/negative helicity components of $g_{\alpha(r)ab}$,

$$\begin{aligned} g_{\alpha(r)ab}^{ab}(\vec{x}) &:= \int \frac{d^3k}{(2\pi)^{\frac{3}{2}}} e^{i\vec{k}\cdot\vec{x}} k X_{\alpha(r)}^{ab}(\vec{k}) \\ &= \oint_{\alpha^b} dt \frac{\dot{\alpha}^b}{(2\pi)^{\frac{3}{2}}} I(\vec{x} - \vec{\alpha}^b(t)). \end{aligned} \quad (\text{B2})$$

$I(\vec{y})$ is defined as

$$\begin{aligned} I(\vec{y}) &:= \int \frac{d^3k}{(2\pi)^{\frac{3}{2}}} e^{i\vec{k}\cdot\vec{y}} k e^{\frac{-k^2 r^2}{2}} \\ &= \frac{4\pi}{(2\pi)^{\frac{3}{2}} r^3 y} \int_0^\infty du \sin\left(\frac{y}{r}u\right) u^2 e^{\frac{-u^2}{2}} \end{aligned} \quad (\text{B3})$$

where we have denoted $|\vec{y}|$ by y .

From (4.6),(3.3) and (B1) it follows that for $|\psi_\Delta\rangle = |\Delta\rangle$ or $|\psi_\Delta\rangle = |\Delta\{\vec{x}_k\}\rangle$

$$\begin{aligned} |\prod_{j=0}^{n-1} \hat{G}_{\alpha_{n-j}}^\dagger \Psi_{weave}[|\psi_\Delta\rangle]| &\leq O^\infty(\epsilon) (\prod_{i=1}^n (\int d^3x_i |G_{\alpha_i(r)}^{*a_i b_i}(\vec{x}_i) G_{\alpha_i(r) a_i b_i}(\vec{x}_i)|^{\frac{1}{2}})) \\ &\leq O^\infty(\epsilon) \prod_{i=1}^n (\int d^3x_i \sum_{a_i, b_i=1..3} (1+4\gamma^2)^{\frac{1}{2}} |g_{\alpha_i(r) a_i b_i}(\vec{x}_i)|). \end{aligned} \quad (\text{B4})$$

We estimate a bound on $|g_{\alpha(r)ab}(\vec{x})|$ through an analysis of its asymptotic behaviour as follows. From [24] we have that

$$\int_0^\infty du \sin(\frac{y}{r}u) u^2 e^{\frac{-u^2}{2}} = \frac{2y}{r} e^{\frac{-y^2}{2r^2}} {}_1F_1(-\frac{1}{2}; \frac{3}{2}; \frac{y^2}{2r^2}), \quad (\text{B5})$$

where ${}_1F_1(a; c; z)$ is the confluent hypergeometric function. From [25] the asymptotic behaviour of this function as $z \rightarrow \infty$ is

$${}_1F_1(a; c; z) = \frac{\Gamma(c)}{\Gamma(a)} e^z z^{a-c} (1 + O(\frac{1}{z})). \quad (\text{B6})$$

The parameters z, a, c in (B6) are identified through (B5),(B2) and (B3) as $z = \frac{|\vec{x} - \vec{\alpha}_b(t)|^2}{2r^2}$, $a = -\frac{1}{2}$ and $c = \frac{3}{2}$. Using (B6) in (B2) and (B3), it is straightforward to obtain the bound

$$|g_{\alpha(r)ab}(\vec{x})| < \frac{2s}{\pi^2 \min_t |\vec{x} - \vec{\alpha}^b(t)|^4} \quad (\text{B7})$$

where $\min_t f(t)$ refers to the minimum value of the function $f(t)$ over the entire loop α_b and the bound is valid for all $\frac{|\vec{x} - \vec{\alpha}^b(t)|}{r} > \lambda$ for sufficiently large λ . Since the triplets of loops $\alpha_i, i = 1..n$ are all confined to a region of size Ss about the origin (see section 3A(i)), we have that $\min_t |\vec{x} - \vec{\alpha}^b(t)| > |\vec{x}| - Ss$.

Then it follows from (B7) that for $|\vec{x}| > r\lambda + 2Ss$ that

$$|g_{\alpha(r)ab}(\vec{x})| < \frac{32s}{\pi^2 |\vec{x}|^4}. \quad (\text{B8})$$

Also note that from (B2) and (B3) the following bound is easily obtained:

$$\int_{|\vec{x}| < r\lambda + 2Ss} d^3x |g_{\alpha(r)ab}(\vec{x})| \leq (\lambda + \frac{2Ss}{r})^3 \frac{s}{\pi r}. \quad (\text{B9})$$

From (B8),(B9) it follows that

$$\sum_{a,b=1..3} \int d^3x |g_{\alpha(r)ab}(\vec{x})| \leq \frac{9s}{\pi r} \{(\lambda + \frac{2Ss}{r})^3 + \frac{128}{(\lambda + \frac{2Ss}{r})}\}. \quad (\text{B10})$$

Using the above bound in (B4) with either (a) or (b) of Appendix A and the ϵ -independence of S, λ, n , we obtain the desired result, namely that

$$\left| \prod_{j=0}^{n-1} \hat{G}_{\alpha_{n-j}}^{\dagger} \Psi_{weave}[\psi_{\Delta} >] \right| \leq O^{\infty}(\epsilon). \quad (\text{B11})$$

APPENDIX C

Let

$$O_I^H(A) \in \left\{ \frac{\text{Tr} H_{\eta}}{2}, \text{Tr} H_{\eta} \tau^k, \eta \in G_{\Delta loop}^s, k = 1..3 \right\}, \quad I = 1..N. \quad (\text{C1})$$

Thus each $O_I^H(A)$ is (half) the trace of a holonomy or the trace of the product of a holonomy with $\tau^i, i \in \{1, 2, 3\}$, the holonomy being around any loop of length at most equal to s . Then the following bounds hold.

$$\left| \left(\prod_{j=0}^{n-1} \int d^3 k_{n-j} (G_{\alpha_{n-j}(r) a_{n-j} b_{n-j}}(\vec{k}_{n-j}))^* \hat{E}_{(r)}^{a_{n-j} b_{n-j}}(\vec{k}_{n-j}) \prod_{I=1}^N O_I^H(A) \right)_{A=0} \right| :=$$

$$\left| \left(\int d^3 k_n (G_{\alpha_n(r) a_n b_n}(\vec{k}_n))^* \hat{E}_{(r)}^{a_n b_n}(\vec{k}_n) \dots \int d^3 k_1 (G_{\alpha_1(r) a_1 b_1}(\vec{k}_1))^* \hat{E}_{(r)}^{a_1 b_1}(\vec{k}_1) \prod_{I=1}^N O_I^H(A) \right)_{A=0} \right|$$

$$\leq N^n \epsilon^{2n} c_n \quad (\text{C2})$$

with $c_n = \left(\frac{27\sqrt{1+4\gamma^2\gamma}}{4\pi^2} \right)^n$.

$$\left| \int \frac{d^3 k}{2i\gamma l_P^2} k[(1 \mp 2i\gamma) \hat{E}_{(r)}^{\pm}(\vec{k}) \hat{E}_{(r)}^{\pm\dagger}(\vec{k})] \right.$$

$$\left. \prod_{j=0}^{n-1} \int d^3 k_{n-j} (G_{\alpha_{n-j}(r) a_{n-j} b_{n-j}}(\vec{k}_{n-j}))^* \hat{E}_{(r)}^{a_{n-j} b_{n-j}}(\vec{k}_{n-j}) \prod_{I=1}^N O_I^H(A) \right|_{A=0}$$

$$\leq N^{n+2} \epsilon^{2n+2} d_n, \quad d_n = \frac{3}{\sqrt{2}} \left(\frac{27\sqrt{1+4\gamma^2\gamma}}{4\pi^2} \right)^{n+1}. \quad (\text{C3})$$

Here $\alpha_i, i = 1..n$ are n triplets of loops, each loop with length at most equal to s , and $\hat{E}^{\pm}$ are the positive and negative helicity components of the triad operator.

Sketch of the proof: We shall only describe the main steps of the proof. The details are straightforward to work out and the interested reader may easily do so.

(i) The following bound holds

$$\left| \left(\prod_{j=0}^{n-1} \hat{E}_{(r)}^{a_{n-j} b_{n-j}}(\vec{k}_{n-j}) O_I^H(A) \right)_{A=0} \right| \leq \left(\frac{l_P^2 \gamma s}{(2\pi)^{\frac{3}{2}}} \right)^n \prod_{i=1}^n e^{\frac{-k_i^2 r^2}{2}}. \quad (\text{C4})$$

Note that the action of the smeared triad operator $\hat{E}_{(r)i}^a(\vec{k})$ on a cylindrical function ψ_α based on a graph α with edges e_J is

$$\hat{E}_{(r)i}^a(\vec{k})\psi_\alpha = il_P^2\gamma \sum_J \int dt \dot{e}_J^a \frac{e^{-i\vec{k}\cdot\vec{e}_J(t)} e^{-\frac{k^2 r^2}{2}}}{(2\pi)^{\frac{3}{2}}} \{ (h_{e_J}(1, t) \tau_i h_{e_J}(t, 0))_A^B \frac{\partial \psi_\alpha}{\partial h_{e_J}(1, 0)_A^B} \}. \quad (C5)$$

For ψ_α of the type O_I^H , the term in curly brackets corresponds to an insertion of τ^i at parameter value t . The action of a product of n factors of triad operators on O_I^H yields n integrations over the loop labelling O_I^H , with n insertions of τ matrices along the loop. The evaluation at $A = 0$ of $O_I^H(A)$ modified by these insertions yields the trace of a product of τ matrices. This trace is bounded by unity. The n integrations along the loop are responsible for the s^n factor in the bound (C4) and the origin of the $(\frac{l_P^2\gamma}{(2\pi)^{\frac{3}{2}}})e^{-\frac{k_i^2 r^2}{2}}$ factors is obvious from (C5).

(ii) The following bound holds.

$$\left| \left(\prod_{j=0}^{n-1} \hat{E}_{(r)}^{a_{n-j} b_{n-j}}(\vec{k}_{n-j}) \prod_{I=1}^N O_I(A) \right)_{A=0} \right| \leq N^n \left(\frac{l_P^2 \gamma s}{(2\pi)^{\frac{3}{2}}} \right)^n \prod_{i=1}^n e^{-\frac{k_i^2 r^2}{2}}. \quad (C6)$$

This follows from a straightforward estimate of the number of terms of type (C4).

(iii) For any ‘smearing’ function $G_{a_1..a_n b_1..b_n}(\vec{k}_1, \dots, \vec{k}_n)$ the following bound holds.

$$\begin{aligned} & \left| \left(\int \prod_{i=1}^n d^3 k_i (G_{a_1..a_n b_1..b_n}(\vec{k}_1, \dots, \vec{k}_n))^* \prod_{j=0}^{n-1} \hat{E}_{(r)}^{a_{n-j} b_{n-j}}(\vec{k}_{n-j}) \prod_{I=1}^N O_I(A) \right)_{A=0} \right| \\ & \leq N^n \left(\frac{9l_P^2 \gamma s}{(2\pi)^{\frac{3}{2}}} \right)^n \int \prod_{i=1}^n d^3 k_i e^{-\frac{k_i^2 r^2}{2}} (\sum_{\{a_i, b_i\}} |G_{a_1..a_n b_1..b_n}(\vec{k}_1, \dots, \vec{k}_n)|^2)^{\frac{1}{2}}. \end{aligned} \quad (C7)$$

To see this, note that the left hand side of the above equation is bounded by the integral of the product of the norms of the smearing function and the quantity $\left(\prod_{j=0}^{n-1} \hat{E}_{(r)}^{a_{n-j} b_{n-j}}(\vec{k}_{n-j}) \prod_{I=1}^N O_I(A) \right)_{A=0}$. The norm of the latter is bounded by the right hand side of (C6) augmented by a factor of 9^n which comes from the sum over the indices a_i, b_i . As a result, equation (C7) follows.

(iv) Set $G_{a_1..a_n b_1..b_n}(\vec{k}_1, \dots, \vec{k}_n) = \prod_{i=1}^n G_{\alpha_i(r) a_i b_i}(\vec{k}_i)$ in (C7). Using (2.11), it is easily verified that

$$\left(\sum_{a,b} |G_{\alpha_i(r) ab}(\vec{k}_i)|^2 \right)^{\frac{1}{2}} \leq k_i e^{-\frac{k_i^2 r^2}{2}} (1 + 4\gamma^2)^{\frac{1}{2}} \frac{3s}{(2\pi)^{\frac{3}{2}}}. \quad (C8)$$

The integral over each $\vec{k}_i$ in (C7) furnishes a factor of $\frac{2\pi}{r^4}$. This, together with $\epsilon = \frac{l_P s}{r^2}$ from (2.21) yields (C2). Similarly, for an appropriate choice of $G_{a_1..a_n b_1..b_n}(\vec{k}_1, \dots, \vec{k}_n)$ and straightforwardly derived bounds thereon, equation (C3) may also be obtained.

APPENDIX D

We prove the identities (5.3) through the following steps.

(1) Consider a set of operators $\hat{O}_i, i = 0..n$ such that each $\hat{O}_i$ is of the type defined in equation (4.9). Denote the operators $\hat{E}_{(r)i_j}^{a_j}$ by $\hat{E}_j$ and the Kronecker deltas $\delta_{i_j}^{a_j}$ by δ_j . Define $\hat{O}_{1,2,..,n}$ and $\hat{O}_{1,2,..,\not{i},\not{j}..,n}$ by

$$\hat{O}_{1,2,..,n} := \hat{O}_1 \hat{O}_2 \dots \hat{O}_n \quad (\text{D1})$$

$$\hat{O}_{1,2,..,\not{i},\not{j}..,n} := \hat{O}_1 \hat{O}_2 \dots \hat{O}_{i-1} [\hat{E}_i, \hat{O}_i \hat{O}_{i+1} \dots \hat{O}_{j-1} [\hat{E}_j \hat{O}_j \dots \hat{O}_n] \dots]. \quad (\text{D2})$$

Thus, in definition (D2), each ‘slashed’ index implies a commutator of the string of operators which follow the index with the triad operator labelled by that index. Then the following lemma holds.

Lemma 3:

$$\begin{aligned} \prod_{i=1}^n (\hat{E}_i - \delta_i) \hat{O}_i &= (E_1 - \delta_1) \hat{O}_1 (E_2 - \delta_2) \hat{O}_2 \dots (E_n - \delta_n) \hat{O}_n \\ &= \hat{O}_{1..n} \prod_{i=1}^n (E_i - \delta_i) + \sum_{I=1}^n \left(\sum_{i_1, i_2, \dots, i_I} \hat{O}_{1,2,..,\not{i}_1,\not{i}_2,..,\not{i}_I..,n} \prod_{j \notin \{i_1, \dots, i_I\}} (E_j - \delta_j) \right), \quad (\text{D3}) \end{aligned}$$

where the second sum in the second term on the right hand side is over all choices of ‘slashed indices’ $i_1 < i_2 \dots < i_I$.

Proof (by induction): Equation (D3) is easily verified for $n = 1$. Let it be true for $n = m$. For $n = m + 1$, the left hand side of (D3) is $\prod_{i=1}^{m+1} (\hat{E}_i - \delta_i) \hat{O}_i$. We apply (D3) for $n = m$ to

the string of operators $\prod_{i=2}^{m+1}(\hat{E}_i - \delta_i)\hat{O}_i$ to obtain

$$\begin{aligned} \prod_{i=1}^{m+1}(\hat{E}_i - \delta_i)\hat{O}_i &= \{(\hat{E}_1 - \delta_1)\hat{O}_{1\dots m+1} \prod_{i=2}^{m+1}(\hat{E}_i - \delta_i)\} \\ &+ \{(\hat{E}_1 - \delta_1)\hat{O}_1 \sum_{I=2}^{m+1} \sum_{i_2, \dots, i_I \neq 1} \hat{O}_{2,3,\dots, \cancel{i_2}, \dots, \cancel{i_I}, \dots, m+1} \prod_{j \notin \{1, i_2, \dots, i_I\}} (\hat{E}_j - \delta_j)\} \end{aligned} \quad (D4)$$

$$\begin{aligned} &= \{\hat{O}_{1\dots m+1} \prod_{i=1}^{m+1}(\hat{E}_i - \delta_i) + \hat{O}_{\cancel{1}, 2, \dots, m+1} \prod_{i=2}^{m+1}(\hat{E}_i - \delta_i)\} \\ &+ \{\sum_{I=2}^{m+1} \sum_{i_2, \dots, i_I \neq 1} \hat{O}_{\cancel{1}, 2, 3, \dots, \cancel{i_2}, \dots, \cancel{i_I}, \dots, m+1} \prod_{j \notin \{1, i_2, \dots, i_I\}} (\hat{E}_j - \delta_j) \\ &+ \sum_{I=2}^{m+1} \sum_{i_2, \dots, i_I \neq 1} \hat{O}_{1, 2, 3, \dots, \cancel{i_2}, \dots, \cancel{i_I}, \dots, m+1} (E_1 - \delta_1) \prod_{j \notin \{1, i_2, \dots, i_I\}} (\hat{E}_j - \delta_j)\} \end{aligned} \quad (D5)$$

$$= \hat{O}_{1\dots m+1} \prod_{i=1}^{m+1}(\hat{E}_i - \delta_i) + \sum_{I=1}^{m+1} \sum_{i_1, i_2, \dots, i_I} \hat{O}_{1, 2, \dots, \cancel{i_1}, \dots, \cancel{i_I}, \dots, m+1} \prod_{j \notin \{i_1, \dots, i_I\}} (E_j - \delta_j) \quad (D6)$$

This completes the proof.

Note: In the equations above it is understood that any set of indices $\{i_k\}$ is such that if $p < q$, then $i_p < i_q$ and that the set of indices range over all appropriate subsets of $\{1, 2, \dots, m+1\}$ modulo explicitly forbidden values (for e.g. in some cases the indices have to be all different from 1).

(2) In the connection representation the commutator $[\hat{E}_{i_1}, \hat{O}_{i_1, i_1+1, \dots, \cancel{j_1}, \dots, \cancel{j_2}, \dots, n}]$ is a ‘multiplication’ operator whose action on any state $\psi(\bar{A})$ is given by

$$\begin{aligned} [\hat{E}_{i_1}, \hat{O}_{i_1, i_1+1, \dots, \cancel{j_1}, \dots, \cancel{j_2}, \dots, n}] \psi(\bar{A}) &= \int d^3y \frac{e^{-\frac{|\vec{x}_{i_1} - \vec{y}|^2}{2r^2}}}{(2\pi r^2)^{\frac{3}{2}}} \\ &(-i l_P^2 \gamma \frac{\delta}{\delta A_a^i(\vec{y})} \hat{O}_{i_1, i_1+1, \dots, \cancel{j_1}, \dots, \cancel{j_2}, \dots, n}(A)|_{A=\bar{A}}) \psi(\bar{A}) \end{aligned} \quad (D7)$$

with the term multiplying $\psi(\bar{A})$ on the right hand side of the above equation completely determined by the holonomies of the generalised connection $\bar{A}$ along graphs contained in G_Δ (recall that G_Δ is the set of all graphs which intersect Δ in at most a finite number of points).

Using (D7) and Lemma 3 above, in conjunction with (a)-(c) of section 4C, it is straight-

forward to see that for $|\psi_\Delta\rangle = |\Delta\rangle$ or $|\psi_\Delta\rangle = |\Delta\{x_k\}\rangle$, we have

$$\begin{aligned} \Psi_0^\epsilon[\hat{O}_0(E_1 - \delta_1)\hat{O}_1(E_2 - \delta_2)\hat{O}_2\cdots(E_n - \delta_n)\hat{O}_n|\psi_\Delta\rangle] &= \Psi_{weave}[|\psi_\Delta\rangle]\Phi_0^\epsilon[\hat{O}_0\hat{O}_{1,2,\dots,\hat{y}_1,\hat{y}_2,\dots,n}|\cdot\rangle] \\ &+ \sum_{I=1}^{n-1}(\sum_{i_1,i_2,\dots,i_I}\Psi_{weave}[\prod_{j\notin\{i_1,\dots,i_I\}}(E_j - \delta_j)|\psi_\Delta\rangle]\Phi_0^\epsilon[\hat{O}_0\hat{O}_{1,2,\dots,\hat{y}_1,\hat{y}_2,\dots,n}|\cdot\rangle] \\ &+ \Psi_{weave}[\prod_{i=1}^n(E_i - \delta_i)|\psi_\Delta\rangle]\Phi_0^\epsilon[\hat{O}_0\hat{O}_{1..n}|\cdot\rangle] \end{aligned} \quad (D8)$$

Note that

$$\hat{O}_0\hat{O}_{1,2,\dots,\hat{y}_1,\hat{y}_2,\dots,n}|\cdot\rangle = \hat{O}_0\hat{O}_1\hat{O}_2\cdots\hat{E}_{i_1}\hat{O}_{i_1}\cdots\hat{E}_{i_I}\hat{O}_{i_I}\cdots\hat{O}_n|\cdot\rangle. \quad (D9)$$

Using this in (D8) and integrating against $\prod_{i=1}^n G_{\alpha_i(r)a_i b_i}^*$ with α_i being triplets of loops in $G_{\Delta loop}^s$, we obtain

$$\begin{aligned} &\Psi_0^\epsilon[\hat{O}_0\hat{G}_{\alpha_1(r)}^\dagger\hat{O}_1\cdots\hat{G}_{\alpha_n(r)}^\dagger\hat{O}_n|\psi_\Delta\rangle] = \\ &\Psi_{weave}[|\psi_\Delta\rangle]\Phi_0^\epsilon[\hat{O}_0(\int G_{\alpha_1(r)a_1 b_1}^*\hat{E}_{(r)}^{a_1 b_1})\hat{O}_1\cdots(\int G_{\alpha_n(r)a_n b_n}^*\hat{E}_{(r)}^{a_n b_n})\hat{O}_n|\cdot\rangle] \\ &+ \sum_{I=1}^{n-1}(\sum_{i_1,i_2,\dots,i_I}\Psi_{weave}[\prod_{j\notin\{i_1,\dots,i_I\}}\hat{G}_{\alpha_{i_j}(r)}^\dagger|\psi_\Delta\rangle] \\ &\Phi_0^\epsilon[\hat{O}_0\hat{O}_1\cdots(\int G_{\alpha_{i_1}(r)a_{i_1} b_{i_1}}^*\hat{E}_{(r)}^{a_{i_1} b_{i_1}})\hat{O}_{i_1}\cdots(\int G_{\alpha_{i_I}(r)a_{i_I} b_{i_I}}^*\hat{E}_{(r)}^{a_{i_I} b_{i_I}})\hat{O}_{i_I}\cdots\hat{O}_n|\cdot\rangle]) \\ &+ \Psi_{weave}[\prod_{i=1}^n\hat{G}_{\alpha_i(r)}^\dagger|\psi_\Delta\rangle]\Phi_0^\epsilon[\hat{O}_0\hat{O}_{1..n}|\cdot\rangle] \end{aligned} \quad (D10)$$

(3) Set

$$\hat{O}_i = \prod_{I=1}^{P_i} \hat{O}_{i_I}^H \text{ for } i = 0..n-1, \quad \hat{O}_n = \hat{O}'_n \prod_{I=1}^N \hat{O}_I^H \quad \hat{O}'_n = \prod_{I=1}^{P_n} \hat{O}_{n_I}^H. \quad (D11)$$

where $\hat{O}_I^H, \hat{O}_{i_I}^H$ are quantum operators corresponding to classical functions of the type defined by (C1). Define $P := \sum_{i=0}^n P_i$ and restrict attention to n, P independent of ϵ . Using (D11), (4.7), (4.7) and (C2), (C3) in conjunction with straightforward bounds on the number of terms of type (C2), (C3), we obtain

$$\begin{aligned} &|\sum_{I=1}^{n-1}(\sum_{i_1,i_2,\dots,i_I}\Psi_{weave}[\prod_{j\notin\{i_1,\dots,i_I\}}\hat{G}_{\alpha_{i_j}(r)}^\dagger|\psi_\Delta\rangle] \\ &\Phi_0^\epsilon[\hat{O}_0\hat{O}_1\cdots(\int G_{\alpha_{i_1}(r)a_{i_1} b_{i_1}}^*\hat{E}_{(r)}^{a_{i_1} b_{i_1}})\hat{O}_{i_1}\cdots(\int G_{\alpha_{i_I}(r)a_{i_I} b_{i_I}}^*\hat{E}_{(r)}^{a_{i_I} b_{i_I}})\hat{O}_{i_I}\cdots\hat{O}_n|\cdot\rangle])| \\ &\leq O^\infty(\epsilon)(N+P)^{n+1}, \end{aligned} \quad (D12)$$

$$|\Psi_{weave}[\prod_{i=1}^n\hat{G}_{\alpha_i(r)}^\dagger|\psi_\Delta\rangle]\Phi_0^\epsilon[\hat{O}_0\hat{O}_{1..n}|\cdot\rangle]| \leq O^\infty(\epsilon)(1+(N+P)^2O(\epsilon^2)). \quad (D13)$$

In the above equations the $O^\infty(\epsilon)$, $O(\epsilon^2)$ terms are independent of N . Thus (D12) and (D13) display the N dependence of the bounds explicitly. Note that since $P_i \geq 1$, we have that $P > n$. Using this in conjunction with the above bounds in (D10) we obtain

$$\begin{aligned} & \Psi_0^\epsilon [\hat{O}_0 \hat{G}_{\alpha_1(r)}^\dagger \hat{O}_1 \dots \hat{G}_{\alpha_n(r)}^\dagger \hat{O}'_n \prod_{I=1}^N \hat{O}_I^H |\psi_\Delta > \\ &= \Psi_{weave} [|\psi_\Delta >] \Phi_0^\epsilon [\hat{O}_0 (\int G_{\alpha_1(r)a_1b_1}^* \hat{E}_{(r)}^{a_1b_1}) \hat{O}_1 \dots (\int G_{\alpha_n(r)a_nb_n}^* \hat{E}_{(r)}^{a_nb_n}) \hat{O}'_n \prod_{I=1}^N \hat{O}_I^H |\cdot >] \\ & \quad + O^{(\infty)}(\epsilon)(N+P)^{n+1}. \end{aligned} \quad (D14)$$

Using the definition of the seminorm in section 4B, it is straightforward to see that (D14) implies that

$$\|\hat{O}'_n \hat{G}_{\alpha_n(r)} \dots \hat{O}_1 \hat{G}_{\alpha_1(r)} \hat{O}_0 \Psi_0^\epsilon\|^\epsilon = \|\Psi^\epsilon\|^\epsilon + O^\infty(\epsilon) \quad (D15)$$

where Ψ^ϵ is a distribution of the type defined in section 4C for which

$$\Phi^\epsilon = \hat{O}'_n (\int G_{\alpha_n(r)a_nb_n} \hat{E}_{(r)}^{a_nb_n}) \dots \hat{O}_1 (\int G_{\alpha_1(r)a_1b_1} \hat{E}_{(r)}^{a_1b_1}) \hat{O}_0 \Phi_0^\epsilon$$

This completes the proof of (5.3).

APPENDIX E

Let β^i be loops of size r . Let $\alpha^i = (\beta^i)^n$ (i.e. α^i is obtained by going around β^i 'n' times) with $n = \frac{s}{r}$. Then it follows that

$$(\Delta \hat{g}_{\alpha(r)})^2 = l_P^2 (1 + 4\gamma^2) \int d^3 k k e^{-k^2 r^2} (|X_{\alpha}^+(\vec{k})|^2 + |X_{\alpha}^-(\vec{k})|^2) \quad (E1)$$

$$= \frac{s^2}{r^2} l_P^2 (1 + 4\gamma^2) \int d^3 k k e^{-k^2 r^2} (|X_{\beta}^+(\vec{k})|^2 + |X_{\beta}^-(\vec{k})|^2) \quad (E2)$$

$$= \frac{s^2}{r^2} \frac{l_P^2}{r^4} (1 + 4\gamma^2) \int d^3 u u e^{-u^2} (|X_{\beta}^+(\frac{\vec{u}}{r})|^2 + |X_{\beta}^-(\frac{\vec{u}}{r})|^2), \quad (E3)$$

where we have defined the dimensionless variable $\vec{u} = r\vec{k}$. Next, note that

$$X_{\beta^k}^a(\frac{\vec{u}}{r}) = r X_{\beta_1^k}^a(\vec{u}), \quad (E4)$$

where

$$X_{\beta_1^k}^a(\vec{u}) := \frac{1}{(2\pi)^{\frac{3}{2}}} \oint_{\beta_1^k} ds e^{-i\vec{u} \cdot \vec{\beta}_1^k(s)} \dot{\beta}_1^a, \quad (E5)$$

with $\vec{\beta}_1^k(s) := \frac{\vec{\beta}^k(s)}{r}$. Since β^k is a loop of length r , β_1^k is a 'unit' size loop. For generic β_1^k we expect that there exist real a, b and positive definite c , all independent of ϵ such that for $u \in [a, b]$, we have that

$$|X_{\beta_1^k}^\pm(\vec{u})| \geq c. \quad (E6)$$

Using (E4) and (E6), we obtain

$$(\Delta \hat{g}_{\alpha(r)})^2 \geq \epsilon^2 (1 + 4\gamma^2) 2c^2 \int_a^b d^3 u u e^{-u^2} \quad (\text{E7})$$

-
- [1] A. Ashtekar, C. Rovelli and L. Smolin, *Phys. Rev.***D44** (1991), 1740.
 - [2] M. Varadarajan, *Phys.Rev.***D64**, 104003 (2001).
 - [3] M. Varadarajan, *Phys.Rev.***D66**, 024017 (2002).
 - [4] C. Rovelli, *Class.Quant.Grav.***8** 1613 (1991).
 - [5] A. Ashtekar and J. Lewandowski, *Class.Quant.Grav.***14** A55 (1997).
 - [6] T. Thiemann and O. Winkler, *Class.Quant.Grav.***18** 2561 (2001); *Class.Quant.Grav.***18** 4629 (2001).
 - [7] A. Ashtekar and J. Lewandowski, *Class.Quant.Grav.***18** L117 (2001); A. Ashtekar, S. Fairhurst and J. Willis, *Class.Quant.Grav.***20** 1031 (2003).
 - [8] T. Thiemann, ‘Complexifier Coherent States for General Relativity’ e-Print Archive: gr-qc/0206037 .
 - [9] Florian Conrady, ‘Free Vacuum for Loop Quantum Gravity’, e-Print Archive: gr-qc/0409036.
 - [10] A. Ashtekar, C. Rovelli and L. Smolin, *Phys.Rev.Lett.***69** 237 (1992); J. Zwegaard, *Phys.Lett.***B300** 217 (1993); R. Borissov, *Phys.Rev.***D49** 923 (1994); N. Grot and C. Rovelli, *Gen.Rel.Grav.***29** 1039 (1997); M. Arnsdorf, ‘Approximating Connections in Loop Quantum Gravity’, e-Print Archive: gr-qc/9910084; A. Corichi and J.M. Reyes, *Int.J.Mod.Phys.***D10** 325 (2001); T. Tsushima, ‘The expectation value of the metric operator with respect to Gaussian weave state in loop quantum gravity’, e-Print Archive: gr-qc/0212117;
 - [11] L. Bombelli, ‘Statistical Geometry of Random Weave States’ in *Proceedings of 9th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories* (Rome 2000), e-Print Archive: gr-qc/0101080; L. Bombelli, A. Corichi and O. Winkler, ‘Semiclassical Quantum Gravity: Statistics of Combinatorial Riemannian Geometries’ e-Print Archive: gr-qc/0409006 .
 - [12] J. Iwasaki and C. Rovelli, *Int.J.Mod.Phys.***D1**, 533 (1993); J. Iwasaki and C. Rovelli, *Class.Quant.Grav.***11**, 1653 (1994).
 - [13] J. Zegwaard, *Class.Quant.Grav.***10** S273 (1993).
 - [14] J.Fernando Barbero, *Phys.Rev.***D51**, 5507 (1995).
 - [15] A. Ashtekar and J. Lewandowski, *J.Gem.Phys.***17** 191 (1995).
 - [16] D. Marolf and J. Mourão, *Commun. Math. Phys.* ,583 (1995).
 - [17] M. Varadarajan and J.A. Zapata, *Class.Quant.Grav.***17** 4085 (2000).
 - [18] H. Sahlmann, T. Thiemann and O. Winkler, *Nucl.Phys.***B606** 401 (2001); M. Arnsdorf and S. Gupta, *Nucl.Phys.***B577** 529 (2000).
 - [19] Y. Yamasaki, *Measures on infinite dimensional vector spaces* (World Scientific 1985).
 - [20] T. Thiemann, *Class.Quant.Grav.***15** 839 (1998).
 - [21] A. Ashtekar, J. Lewandowski, D. Marolf, J. Mourao, T. Thiemann *J.Math.Phys.***36** 6456 (1995).
 - [22] K. Kuchar, *Phys.Rev.***D4** 55 (1971).
 - [23] See, for example, M.J. Lighthill *Introduction to Fourier analysis and Generalised Functions*, pg 69-70 (Cambridge University Press, 1975).

- [24] *Tables of Integral Transforms Volume I* ed. A. Erdelyi, W. Magnus, F. Oberhettinger and F.C. Tricomi (McGraw-Hill, 1954).
- [25] *Higher Transcendental Functions Volume I* ed. A. Erdelyi, W. Magnus, F. Oberhettinger and F.C. Tricomi (McGraw-Hill, 1954).